



## Target Species

Based on 344,773 complete checklists

**453** species observed in Florida, US (region/US-FL) in Feb that you need for your Florida Life List

|  |                        |
|--|------------------------|
| 1. <b>Black-bellied Whistling-Duck</b> <i>Dendrocygna autumnalis</i> | <b>3.81%</b> frequency |
| 2. <b>Fulvous Whistling-Duck</b> <i>Dendrocygna bicolor</i>          | <b>0.4716%</b>         |
| 3. <b>Snow Goose</b> <i>Anser caerulescens</i>                       | <b>0.1175%</b>         |
| 4. <b>Ross's Goose</b> <i>Anser rossii</i>                           | <b>0.0426%</b>         |
| 5. <b>Greater White-fronted Goose</b> <i>Anser albifrons</i>         | <b>0.0264%</b>         |
| 6. <b>Brant</b> <i>Branta bernicla</i>                               | <b>0.0165%</b>         |
| 7. <b>Cackling Goose</b> <i>Branta hutchinsii</i>                    | <b>0.0006%</b>         |
| 8. <b>Canada Goose</b> <i>Branta canadensis</i>                      | <b>2.58%</b>           |
| 9. <b>Tundra Swan</b> <i>Cygnus columbianus</i>                      | <b>0.0017%</b>         |
| 10. <b>Egyptian Goose</b> <i>Alopochen aegyptiaca</i>                | <b>1.64%</b>           |
| 11. <b>Muscovy Duck</b> <i>Cairina moschata</i>                      | <b>5.03%</b>           |
| 12. <b>Wood Duck</b> <i>Aix sponsa</i>                               | <b>3.42%</b>           |
| 13. <b>Blue-winged Teal</b> <i>Spatula discors</i>                   | <b>9.19%</b>           |
| 14. <b>Cinnamon Teal</b> <i>Spatula cyanoptera</i>                   | <b>0.1734%</b>         |
| 15. <b>Northern Shoveler</b> <i>Spatula clypeata</i>                 | <b>3.00%</b>           |
| 16. <b>Gadwall</b> <i>Mareca strepera</i>                            | <b>0.8058%</b>         |
| 17. <b>Eurasian Wigeon</b> <i>Mareca penelope</i>                    | <b>0.0545%</b>         |

|  |         |
|--|---------|
| 18. <b>American Wigeon</b> <i>Mareca americana</i>         | 1.40%   |
| 19. <b>Mallard</b> <i>Anas platyrhynchos</i>               | 3.63%   |
| 20. <b>American Black Duck</b> <i>Anas rubripes</i>        | 0.0412% |
| 21. <b>Mottled Duck</b> <i>Anas fulvigula</i>              | 9.36%   |
| 22. <b>White-cheeked Pintail</b> <i>Anas bahamensis</i>    | 0.0081% |
| 23. <b>Northern Pintail</b> <i>Anas acuta</i>              | 0.8501% |
| 24. <b>Green-winged Teal</b> <i>Anas crecca</i>            | 1.74%   |
| 25. <b>Canvasback</b> <i>Aythya valisineria</i>            | 0.4040% |
| 26. <b>Redhead</b> <i>Aythya americana</i>                 | 1.33%   |
| 27. <b>Ring-necked Duck</b> <i>Aythya collaris</i>         | 4.20%   |
| 28. <b>Greater Scaup</b> <i>Aythya marila</i>              | 0.4693% |
| 29. <b>Lesser Scaup</b> <i>Aythya affinis</i>              | 5.04%   |
| 30. <b>King Eider</b> <i>Somateria spectabilis</i>         | 0.0000% |
| 31. <b>Common Eider</b> <i>Somateria mollissima</i>        | 0.0452% |
| 32. <b>Harlequin Duck</b> <i>Histrionicus histrionicus</i> | 0.0119% |
| 33. <b>Surf Scoter</b> <i>Melanitta perspicillata</i>      | 0.0841% |
| 34. <b>White-winged Scoter</b> <i>Melanitta deglandi</i>   | 0.0627% |
| 35. <b>Black Scoter</b> <i>Melanitta americana</i>         | 0.2399% |
| 36. <b>Long-tailed Duck</b> <i>Clangula hyemalis</i>       | 0.0403% |
| 37. <b>Bufflehead</b> <i>Bucephala albeola</i>             | 2.09%   |
| 38. <b>Common Goldeneye</b> <i>Bucephala clangula</i>      | 0.2007% |
| 39. <b>Hooded Merganser</b> <i>Lophodytes cucullatus</i>   | 3.93%   |
| 40. <b>Common Merganser</b> <i>Mergus merganser</i>        | 0.0006% |

|   |         |
|---|---------|
| 41. <b>Red-breasted Merganser</b> <i>Mergus serrator</i>        | 3.41%   |
| 42. <b>Masked Duck</b> <i>Nomonyx dominicus</i>                 | 0.0055% |
| 43. <b>Ruddy Duck</b> <i>Oxyura jamaicensis</i>                 | 1.03%   |
| 44. <b>Northern Bobwhite</b> <i>Colinus virginianus</i>         | 0.6601% |
| 45. <b>Wild Turkey</b> <i>Meleagris gallopavo</i>               | 1.65%   |
| 46. <b>American Flamingo</b> <i>Phoenicopterus ruber</i>        | 0.1955% |
| 47. <b>Least Grebe</b> <i>Tachybaptus dominicus</i>             | 0.0093% |
| 48. <b>Pied-billed Grebe</b> <i>Podilymbus podiceps</i>         | 16.31%  |
| 49. <b>Horned Grebe</b> <i>Podiceps auritus</i>                 | 1.05%   |
| 50. <b>Red-necked Grebe</b> <i>Podiceps grisegena</i>           | 0.0032% |
| 51. <b>Eared Grebe</b> <i>Podiceps nigricollis</i>              | 0.0194% |
| 52. <b>Western Grebe</b> <i>Aechmophorus occidentalis</i>       | 0.0003% |
| 53. <b>Rock Pigeon</b> <i>Columba livia</i>                     | 6.36%   |
| 54. <b>White-crowned Pigeon</b> <i>Patagioenas leucocephala</i> | 0.3591% |
| 55. <b>Eurasian Collared-Dove</b> <i>Streptopelia decaocto</i>  | 9.74%   |
| 56. <b>Inca Dove</b> <i>Columbina inca</i>                      | 0.0038% |
| 57. <b>Common Ground Dove</b> <i>Columbina passerina</i>        | 8.01%   |
| 58. <b>Key West Quail-Dove</b> <i>Geotrygon chrysia</i>         | 0.0058% |
| 59. <b>White-winged Dove</b> <i>Zenaida asiatica</i>            | 3.54%   |
| 60. <b>Zenaida Dove</b> <i>Zenaida aurita</i>                   | 0.0229% |
| 61. <b>Mourning Dove</b> <i>Zenaida macroura</i>                | 40.92%  |
| 62. <b>Smooth-billed Ani</b> <i>Crotophaga ani</i>              | 0.0960% |
| 63. <b>Groove-billed Ani</b> <i>Crotophaga sulcirostris</i>     | 0.0418% |

|     |  |                |
|-----|--|----------------|
| 64. | <b>Dark-billed Cuckoo</b> <i>Coccyzus melacoryphus</i>         | <b>0.0304%</b> |
| 65. | <b>Yellow-billed Cuckoo</b> <i>Coccyzus americanus</i>         | <b>0.0012%</b> |
| 66. | <b>Mangrove Cuckoo</b> <i>Coccyzus minor</i>                   | <b>0.0693%</b> |
| 67. | <b>Lesser Nighthawk</b> <i>Chordeiles acutipennis</i>          | <b>0.0212%</b> |
| 68. | <b>Common Nighthawk</b> <i>Chordeiles minor</i>                | <b>0.0014%</b> |
| 69. | <b>Chuck-will's-widow</b> <i>Antrostomus carolinensis</i>      | <b>0.1758%</b> |
| 70. | <b>Eastern Whip-poor-will</b> <i>Antrostomus vociferus</i>     | <b>0.2491%</b> |
| 71. | <b>Chimney Swift</b> <i>Chaetura pelagica</i>                  | <b>0.0006%</b> |
| 72. | <b>Vaux's Swift</b> <i>Chaetura vauxi</i>                      | <b>0.0278%</b> |
| 73. | <b>Ruby-throated Hummingbird</b> <i>Archilochus colubris</i>   | <b>2.83%</b>   |
| 74. | <b>Black-chinned Hummingbird</b> <i>Archilochus alexandri</i>  | <b>0.1015%</b> |
| 75. | <b>Calliope Hummingbird</b> <i>Selasphorus calliope</i>        | <b>0.0302%</b> |
| 76. | <b>Rufous Hummingbird</b> <i>Selasphorus rufus</i>             | <b>0.2042%</b> |
| 77. | <b>Allen's Hummingbird</b> <i>Selasphorus sasin</i>            | <b>0.0061%</b> |
| 78. | <b>Broad-tailed Hummingbird</b> <i>Selasphorus platycercus</i> | <b>0.0000%</b> |
| 79. | <b>Broad-billed Hummingbird</b> <i>Cynanthus latirostris</i>   | <b>0.0220%</b> |
| 80. | <b>Buff-bellied Hummingbird</b> <i>Amazilia yucatanensis</i>   | <b>0.0644%</b> |
| 81. | <b>King Rail</b> <i>Rallus elegans</i>                         | <b>0.4655%</b> |
| 82. | <b>Clapper Rail</b> <i>Rallus crepitans</i>                    | <b>0.9769%</b> |
| 83. | <b>Virginia Rail</b> <i>Rallus limicola</i>                    | <b>0.2663%</b> |
| 84. | <b>Sora</b> <i>Porzana carolina</i>                            | <b>2.10%</b>   |
| 85. | <b>Common Gallinule</b> <i>Gallinula galeata</i>               | <b>19.10%</b>  |

|  |                |
|--|----------------|
| 86. <b>American Coot</b> <i>Fulica americana</i>               | <b>13.06%</b>  |
| 87. <b>Purple Gallinule</b> <i>Porphyrio martinica</i>         | <b>4.18%</b>   |
| 88. <b>Gray-headed Swamphen</b> <i>Porphyrio poliocephalus</i> | <b>1.68%</b>   |
| 89. <b>Yellow Rail</b> <i>Coturnicops noveboracensis</i>       | <b>0.0009%</b> |
| 90. <b>Black Rail</b> <i>Laterallus jamaicensis</i>            | <b>0.0154%</b> |
| 91. <b>Limpkin</b> <i>Aramus guarauna</i>                      | <b>9.27%</b>   |
| 92. <b>Sandhill Crane</b> <i>Antigone canadensis</i>           | <b>11.18%</b>  |
| 93. <b>Whooping Crane</b> <i>Grus americana</i>                | <b>0.1616%</b> |
| 94. <b>Black-necked Stilt</b> <i>Himantopus mexicanus</i>      | <b>0.8603%</b> |
| 95. <b>American Avocet</b> <i>Recurvirostra americana</i>      | <b>0.9203%</b> |
| 96. <b>American Oystercatcher</b> <i>Haematopus palliatus</i>  | <b>1.39%</b>   |
| 97. <b>Black-bellied Plover</b> <i>Pluvialis squatarola</i>    | <b>4.25%</b>   |
| 98. <b>American Golden-Plover</b> <i>Pluvialis dominica</i>    | <b>0.0081%</b> |
| 99. <b>Pacific Golden-Plover</b> <i>Pluvialis fulva</i>        | <b>0.0003%</b> |
| 100. <b>Killdeer</b> <i>Charadrius vociferus</i>               | <b>10.00%</b>  |
| 101. <b>Semipalmated Plover</b> <i>Charadrius semipalmatus</i> | <b>2.48%</b>   |
| 102. <b>Piping Plover</b> <i>Charadrius melodus</i>            | <b>0.8310%</b> |
| 103. <b>Wilson's Plover</b> <i>Anarhynchus wilsonia</i>        | <b>0.8974%</b> |
| 104. <b>Mountain Plover</b> <i>Anarhynchus montanus</i>        | <b>0.0003%</b> |
| 105. <b>Snowy Plover</b> <i>Anarhynchus nivosus</i>            | <b>0.4983%</b> |
| 106. <b>Whimbrel</b> <i>Numenius phaeopus</i>                  | <b>0.1407%</b> |
| 107. <b>Long-billed Curlew</b> <i>Numenius americanus</i>      | <b>0.0597%</b> |

|      |   |                |
|------|---|----------------|
| 108. | <b>Bar-tailed Godwit</b> <i>Limosa lapponica</i>            | <b>0.0032%</b> |
| 109. | <b>Black-tailed Godwit</b> <i>Limosa limosa</i>             | <b>0.0009%</b> |
| 110. | <b>Marbled Godwit</b> <i>Limosa fedoa</i>                   | <b>0.8243%</b> |
| 111. | <b>Short-billed Dowitcher</b> <i>Limnodromus griseus</i>    | <b>1.80%</b>   |
| 112. | <b>Long-billed Dowitcher</b> <i>Limnodromus scolopaceus</i> | <b>1.09%</b>   |
| 113. | <b>American Woodcock</b> <i>Scolopax minor</i>              | <b>0.0928%</b> |
| 114. | <b>Wilson's Snipe</b> <i>Gallinago delicata</i>             | <b>2.99%</b>   |
| 115. | <b>Wilson's Phalarope</b> <i>Phalaropus tricolor</i>        | <b>0.0032%</b> |
| 116. | <b>Red Phalarope</b> <i>Phalaropus fulicarius</i>           | <b>0.0046%</b> |
| 117. | <b>Spotted Sandpiper</b> <i>Actitis macularius</i>          | <b>2.37%</b>   |
| 118. | <b>Solitary Sandpiper</b> <i>Tringa solitaria</i>           | <b>0.1966%</b> |
| 119. | <b>Lesser Yellowlegs</b> <i>Tringa flavipes</i>             | <b>3.06%</b>   |
| 120. | <b>Willet</b> <i>Tringa semipalmata</i>                     | <b>7.08%</b>   |
| 121. | <b>Greater Yellowlegs</b> <i>Tringa melanoleuca</i>         | <b>4.53%</b>   |
| 122. | <b>Ruddy Turnstone</b> <i>Arenaria interpres</i>            | <b>5.47%</b>   |
| 123. | <b>Red Knot</b> <i>Calidris canutus</i>                     | <b>0.8249%</b> |
| 124. | <b>Ruff</b> <i>Calidris pugnax</i>                          | <b>0.0203%</b> |
| 125. | <b>Curlew Sandpiper</b> <i>Calidris ferruginea</i>          | <b>0.0003%</b> |
| 126. | <b>Stilt Sandpiper</b> <i>Calidris himantopus</i>           | <b>0.1966%</b> |
| 127. | <b>Sanderling</b> <i>Calidris alba</i>                      | <b>5.94%</b>   |
| 128. | <b>Dunlin</b> <i>Calidris alpina</i>                        | <b>3.20%</b>   |
| 129. | <b>Purple Sandpiper</b> <i>Calidris maritima</i>            | <b>0.1143%</b> |
| 130. | <b>White-rumped Sandpiper</b> <i>Calidris fuscicollis</i>   | <b>0.0003%</b> |

|  |                |
|--|----------------|
| 131. <b>Least Sandpiper</b> <i>Calidris minutilla</i>            | <b>3.45%</b>   |
| 132. <b>Pectoral Sandpiper</b> <i>Calidris melanotos</i>         | <b>0.0168%</b> |
| 133. <b>Western Sandpiper</b> <i>Calidris mauri</i>              | <b>1.34%</b>   |
| 134. <b>Semipalmated Sandpiper</b> <i>Calidris pusilla</i>       | <b>0.0044%</b> |
| 135. <b>Long-tailed Jaeger</b> <i>Stercorarius longicaudus</i>   | <b>0.0003%</b> |
| 136. <b>Parasitic Jaeger</b> <i>Stercorarius parasiticus</i>     | <b>0.0821%</b> |
| 137. <b>Pomarine Jaeger</b> <i>Stercorarius pomarinus</i>        | <b>0.0395%</b> |
| 138. <b>Atlantic Puffin</b> <i>Fratercula arctica</i>            | <b>0.0003%</b> |
| 139. <b>Razorbill</b> <i>Alca torda</i>                          | <b>0.0067%</b> |
| 140. <b>Dovekie</b> <i>Alle alle</i>                             | <b>0.0000%</b> |
| 141. <b>Thick-billed Murre</b> <i>Uria lomvia</i>                | <b>0.0049%</b> |
| 142. <b>Black-legged Kittiwake</b> <i>Rissa tridactyla</i>       | <b>0.0145%</b> |
| 143. <b>Bonaparte's Gull</b> <i>Chroicocephalus philadelphia</i> | <b>2.59%</b>   |
| 144. <b>Black-headed Gull</b> <i>Chroicocephalus ridibundus</i>  | <b>0.0481%</b> |
| 145. <b>Laughing Gull</b> <i>Leucophaeus atricilla</i>           | <b>16.97%</b>  |
| 146. <b>Franklin's Gull</b> <i>Leucophaeus pipixcan</i>          | <b>0.0096%</b> |
| 147. <b>Black-tailed Gull</b> <i>Larus crassirostris</i>         | <b>0.0003%</b> |
| 148. <b>Heermann's Gull</b> <i>Larus heermanni</i>               | <b>0.0128%</b> |
| 149. <b>Ring-billed Gull</b> <i>Larus delawarensis</i>           | <b>18.90%</b>  |
| 150. <b>Herring Gull</b> <i>Larus argentatus</i>                 | <b>5.05%</b>   |
| 151. <b>Great Black-backed Gull</b> <i>Larus marinus</i>         | <b>1.09%</b>   |
| 152. <b>Glaucous Gull</b> <i>Larus hyperboreus</i>               | <b>0.0444%</b> |
| 153. <b>Lesser Black-backed Gull</b> <i>Larus fuscus</i>         | <b>1.69%</b>   |

|      |   |         |
|------|---|---------|
| 154. | <b>California Gull</b> <i>Larus californicus</i>        | 0.0217% |
| 155. | <b>Slaty-backed Gull</b> <i>Larus schistisagus</i>      | 0.0006% |
| 156. | <b>Iceland Gull</b> <i>Larus glaucooides</i>            | 0.0661% |
| 157. | <b>Black Skimmer</b> <i>Rynchops niger</i>              | 3.71%   |
| 158. | <b>Brown Noddy</b> <i>Anous stolidus</i>                | 0.0281% |
| 159. | <b>Sooty Tern</b> <i>Onychoprion fuscatus</i>           | 0.0366% |
| 160. | <b>Bridled Tern</b> <i>Onychoprion anaethetus</i>       | 0.0006% |
| 161. | <b>Least Tern</b> <i>Sternula antillarum</i>            | 0.0003% |
| 162. | <b>Gull-billed Tern</b> <i>Gelochelidon nilotica</i>    | 0.0830% |
| 163. | <b>Caspian Tern</b> <i>Hydroprogne caspia</i>           | 2.92%   |
| 164. | <b>Black Tern</b> <i>Chlidonias niger</i>               | 0.0012% |
| 165. | <b>Forster's Tern</b> <i>Sterna forsteri</i>            | 7.86%   |
| 166. | <b>Common Tern</b> <i>Sterna hirundo</i>                | 0.0757% |
| 167. | <b>Roseate Tern</b> <i>Sterna dougallii</i>             | 0.0017% |
| 168. | <b>Sandwich Tern</b> <i>Thalasseus sandvicensis</i>     | 1.67%   |
| 169. | <b>Elegant Tern</b> <i>Thalasseus elegans</i>           | 0.0072% |
| 170. | <b>Royal Tern</b> <i>Thalasseus maximus</i>             | 8.94%   |
| 171. | <b>White-tailed Tropicbird</b> <i>Phaethon lepturus</i> | 0.0000% |
| 172. | <b>Red-throated Loon</b> <i>Gavia stellata</i>          | 0.0432% |
| 173. | <b>Pacific Loon</b> <i>Gavia pacifica</i>               | 0.0191% |
| 174. | <b>Common Loon</b> <i>Gavia immer</i>                   | 2.62%   |
| 175. | <b>Black-capped Petrel</b> <i>Pterodroma hasitata</i>   | 0.0003% |
| 176. | <b>Great Shearwater</b> <i>Ardenna gravis</i>           | 0.0023% |



|   |         |
|---|---------|
| 177. <b>Sooty Shearwater</b> <i>Ardenna grisea</i>                  | 0.0009% |
| 178. <b>Manx Shearwater</b> <i>Puffinus puffinus</i>                | 0.0026% |
| 179. <b>Audubon's Shearwater</b> <i>Puffinus lherminieri</i>        | 0.0020% |
| 180. <b>Wood Stork</b> <i>Mycteria americana</i>                    | 14.76%  |
| 181. <b>Magnificent Frigatebird</b> <i>Fregata magnificens</i>      | 1.30%   |
| 182. <b>Northern Gannet</b> <i>Morus bassanus</i>                   | 1.31%   |
| 183. <b>Red-footed Booby</b> <i>Sula sula</i>                       | 0.0009% |
| 184. <b>Brown Booby</b> <i>Sula leucogaster</i>                     | 0.1015% |
| 185. <b>Masked Booby</b> <i>Sula dactylatra</i>                     | 0.0203% |
| 186. <b>Anhinga</b> <i>Anhinga anhinga</i>                          | 26.48%  |
| 187. <b>Great Cormorant</b> <i>Phalacrocorax carbo</i>              | 0.0075% |
| 188. <b>Double-crested Cormorant</b> <i>Nannopterum auritum</i>     | 30.71%  |
| 189. <b>Neotropic Cormorant</b> <i>Nannopterum brasilianum</i>      | 0.0682% |
| 190. <b>American White Pelican</b> <i>Pelecanus erythrorhynchos</i> | 7.61%   |
| 191. <b>Brown Pelican</b> <i>Pelecanus occidentalis</i>             | 18.10%  |
| 192. <b>American Bittern</b> <i>Botaurus lentiginosus</i>           | 1.66%   |
| 193. <b>Least Bittern</b> <i>Ixobrychus exilis</i>                  | 0.5960% |
| 194. <b>Yellow-crowned Night Heron</b> <i>Nyctanassa violacea</i>   | 2.68%   |
| 195. <b>Black-crowned Night Heron</b> <i>Nycticorax nycticorax</i>  | 4.76%   |
| 196. <b>Little Blue Heron</b> <i>Egretta caerulea</i>               | 25.76%  |
| 197. <b>Tricolored Heron</b> <i>Egretta tricolor</i>                | 20.65%  |
| 198. <b>Reddish Egret</b> <i>Egretta rufescens</i>                  | 2.91%   |

|      |   |         |
|------|---|---------|
| 199. | <b>Snowy Egret</b> <i>Egretta thula</i>                 | 21.81%  |
| 200. | <b>Green Heron</b> <i>Butorides virescens</i>           | 8.44%   |
| 201. | <b>Western Cattle Egret</b> <i>Bubulcus ibis</i>        | 12.00%  |
| 202. | <b>Great Egret</b> <i>Ardea alba</i>                    | 34.20%  |
| 203. | <b>Great Blue Heron</b> <i>Ardea herodias</i>           | 30.55%  |
| 204. | <b>White Ibis</b> <i>Eudocimus albus</i>                | 34.84%  |
| 205. | <b>Glossy Ibis</b> <i>Plegadis falcinellus</i>          | 11.73%  |
| 206. | <b>White-faced Ibis</b> <i>Plegadis chihi</i>           | 0.0969% |
| 207. | <b>Roseate Spoonbill</b> <i>Platalea ajaja</i>          | 6.66%   |
| 208. | <b>Black Vulture</b> <i>Coragyps atratus</i>            | 24.83%  |
| 209. | <b>Turkey Vulture</b> <i>Cathartes aura</i>             | 43.84%  |
| 210. | <b>Osprey</b> <i>Pandion haliaetus</i>                  | 31.50%  |
| 211. | <b>White-tailed Kite</b> <i>Elanus leucurus</i>         | 0.1265% |
| 212. | <b>Swallow-tailed Kite</b> <i>Elanoides forficatus</i>  | 1.17%   |
| 213. | <b>Golden Eagle</b> <i>Aquila chrysaetos</i>            | 0.0006% |
| 214. | <b>Snail Kite</b> <i>Rostrhamus sociabilis</i>          | 1.53%   |
| 215. | <b>Mississippi Kite</b> <i>Ictinia mississippiensis</i> | 0.0003% |
| 216. | <b>Northern Harrier</b> <i>Circus hudsonius</i>         | 5.38%   |
| 217. | <b>Sharp-shinned Hawk</b> <i>Accipiter striatus</i>     | 0.9807% |
| 218. | <b>Cooper's Hawk</b> <i>Accipiter cooperii</i>          | 3.03%   |
| 219. | <b>Bald Eagle</b> <i>Haliaeetus leucocephalus</i>       | 12.92%  |
| 220. | <b>Red-shouldered Hawk</b> <i>Buteo lineatus</i>        | 27.16%  |
| 221. | <b>Broad-winged Hawk</b> <i>Buteo platypterus</i>       | 0.2486% |

|      |  |                |
|------|--|----------------|
| 222. | <b>Short-tailed Hawk</b> <i>Buteo brachyurus</i>               | <b>0.9438%</b> |
| 223. | <b>Swainson's Hawk</b> <i>Buteo swainsoni</i>                  | <b>0.0566%</b> |
| 224. | <b>Red-tailed Hawk</b> <i>Buteo jamaicensis</i>                | <b>4.00%</b>   |
| 225. | <b>Rough-legged Hawk</b> <i>Buteo lagopus</i>                  | <b>0.0029%</b> |
| 226. | <b>Barn Owl</b> <i>Tyto alba</i>                               | <b>0.2027%</b> |
| 227. | <b>Eastern Screech-Owl</b> <i>Megascops asio</i>               | <b>0.6053%</b> |
| 228. | <b>Great Horned Owl</b> <i>Bubo virginianus</i>                | <b>1.88%</b>   |
| 229. | <b>Burrowing Owl</b> <i>Athene cunicularia</i>                 | <b>0.5218%</b> |
| 230. | <b>Barred Owl</b> <i>Strix varia</i>                           | <b>3.48%</b>   |
| 231. | <b>Short-eared Owl</b> <i>Asio flammeus</i>                    | <b>0.0447%</b> |
| 232. | <b>Belted Kingfisher</b> <i>Megaceryle alcyon</i>              | <b>15.57%</b>  |
| 233. | <b>Yellow-bellied Sapsucker</b> <i>Sphyrapicus varius</i>      | <b>5.80%</b>   |
| 234. | <b>Red-headed Woodpecker</b> <i>Melanerpes erythrocephalus</i> | <b>1.55%</b>   |
| 235. | <b>Red-bellied Woodpecker</b> <i>Melanerpes carolinus</i>      | <b>39.56%</b>  |
| 236. | <b>Downy Woodpecker</b> <i>Dryobates pubescens</i>             | <b>17.92%</b>  |
| 237. | <b>Red-cockaded Woodpecker</b> <i>Dryobates borealis</i>       | <b>0.4336%</b> |
| 238. | <b>Hairy Woodpecker</b> <i>Dryobates villosus</i>              | <b>0.2590%</b> |
| 239. | <b>Pileated Woodpecker</b> <i>Dryocopus pileatus</i>           | <b>13.26%</b>  |
| 240. | <b>Northern Flicker</b> <i>Colaptes auratus</i>                | <b>4.04%</b>   |
| 241. | <b>Crested Caracara</b> <i>Caracara plancus</i>                | <b>1.16%</b>   |
| 242. | <b>Yellow-headed Caracara</b> <i>Daptrius chimachima</i>       | <b>0.0072%</b> |
| 243. | <b>Eurasian Kestrel</b> <i>Falco tinnunculus</i>               | <b>0.0012%</b> |

|      |  |                |
|------|--|----------------|
| 244. | <b>American Kestrel</b> <i>Falco sparverius</i>                | <b>10.64%</b>  |
| 245. | <b>Merlin</b> <i>Falco columbarius</i>                         | <b>0.9235%</b> |
| 246. | <b>Peregrine Falcon</b> <i>Falco peregrinus</i>                | <b>0.8063%</b> |
| 247. | <b>Budgerigar</b> <i>Melopsittacus undulatus</i>               | <b>0.0046%</b> |
| 248. | <b>Monk Parakeet</b> <i>Myiopsitta monachus</i>                | <b>1.68%</b>   |
| 249. | <b>White-winged Parakeet</b> <i>Brotogeris versicolurus</i>    | <b>0.0247%</b> |
| 250. | <b>Carolina Parakeet</b> <i>Conuropsis carolinensis</i>        | <b>0.0003%</b> |
| 251. | <b>Nanday Parakeet</b> <i>Aratinga nenday</i>                  | <b>2.55%</b>   |
| 252. | <b>Mitred Parakeet</b> <i>Psittacara mitratus</i>              | <b>0.1937%</b> |
| 253. | <b>Cuban Pewee</b> <i>Contopus caribaeus</i>                   | <b>0.0099%</b> |
| 254. | <b>Yellow-bellied Flycatcher</b> <i>Empidonax flaviventris</i> | <b>0.0014%</b> |
| 255. | <b>Willow Flycatcher</b> <i>Empidonax traillii</i>             | <b>0.0102%</b> |
| 256. | <b>Least Flycatcher</b> <i>Empidonax minimus</i>               | <b>0.1166%</b> |
| 257. | <b>Hammond's Flycatcher</b> <i>Empidonax hammondii</i>         | <b>0.0215%</b> |
| 258. | <b>Western Flycatcher</b> <i>Empidonax difficilis</i>          | <b>0.0046%</b> |
| 259. | <b>Eastern Phoebe</b> <i>Sayornis phoebe</i>                   | <b>21.43%</b>  |
| 260. | <b>Say's Phoebe</b> <i>Sayornis saya</i>                       | <b>0.0209%</b> |
| 261. | <b>Vermilion Flycatcher</b> <i>Pyrocephalus rubinus</i>        | <b>0.4191%</b> |
| 262. | <b>Ash-throated Flycatcher</b> <i>Myiarchus cinerascens</i>    | <b>0.1752%</b> |
| 263. | <b>Great Crested Flycatcher</b> <i>Myiarchus crinitus</i>      | <b>3.12%</b>   |
| 264. | <b>Brown-crested Flycatcher</b> <i>Myiarchus tyrannulus</i>    | <b>0.0728%</b> |
| 265. | <b>La Sagra's Flycatcher</b> <i>Myiarchus sagrae</i>           | <b>0.0711%</b> |

|      |   |         |
|------|---|---------|
| 266. | <b>Tropical Kingbird</b> <i>Tyrannus melancholicus</i>      | 0.1703% |
| 267. | <b>Couch's Kingbird</b> <i>Tyrannus couchii</i>             | 0.0075% |
| 268. | <b>Cassin's Kingbird</b> <i>Tyrannus vociferans</i>         | 0.0502% |
| 269. | <b>Western Kingbird</b> <i>Tyrannus verticalis</i>          | 0.4644% |
| 270. | <b>Eastern Kingbird</b> <i>Tyrannus tyrannus</i>            | 0.0003% |
| 271. | <b>Gray Kingbird</b> <i>Tyrannus dominicensis</i>           | 0.0238% |
| 272. | <b>Scissor-tailed Flycatcher</b> <i>Tyrannus forficatus</i> | 0.3237% |
| 273. | <b>Fork-tailed Flycatcher</b> <i>Tyrannus savana</i>        | 0.0038% |
| 274. | <b>White-eyed Vireo</b> <i>Vireo griseus</i>                | 10.41%  |
| 275. | <b>Thick-billed Vireo</b> <i>Vireo crassirostris</i>        | 0.0435% |
| 276. | <b>Bell's Vireo</b> <i>Vireo bellii</i>                     | 0.0421% |
| 277. | <b>Yellow-throated Vireo</b> <i>Vireo flavifrons</i>        | 0.2752% |
| 278. | <b>Blue-headed Vireo</b> <i>Vireo solitarius</i>            | 4.92%   |
| 279. | <b>Philadelphia Vireo</b> <i>Vireo philadelphicus</i>       | 0.0078% |
| 280. | <b>Warbling Vireo</b> <i>Vireo gilvus</i>                   | 0.0090% |
| 281. | <b>Red-eyed Vireo</b> <i>Vireo olivaceus</i>                | 0.0012% |
| 282. | <b>Black-whiskered Vireo</b> <i>Vireo altiloquus</i>        | 0.0006% |
| 283. | <b>Loggerhead Shrike</b> <i>Lanius ludovicianus</i>         | 7.99%   |
| 284. | <b>Blue Jay</b> <i>Cyanocitta cristata</i>                  | 33.77%  |
| 285. | <b>Florida Scrub-Jay</b> <i>Aphelocoma coerulescens</i>     | 1.26%   |
| 286. | <b>American Crow</b> <i>Corvus brachyrhynchos</i>           | 18.87%  |
| 287. | <b>Fish Crow</b> <i>Corvus ossifragus</i>                   | 24.81%  |
| 288. | <b>Carolina Chickadee</b> <i>Poecile carolinensis</i>       | 11.49%  |

|      |  |                |
|------|--|----------------|
| 289. | <b>Tufted Titmouse</b> <i>Baeolophus bicolor</i>                       | <b>19.88%</b>  |
| 290. | <b>Horned Lark</b> <i>Eremophila alpestris</i>                         | <b>0.0084%</b> |
| 291. | <b>Bank Swallow</b> <i>Riparia riparia</i>                             | <b>0.0046%</b> |
| 292. | <b>Tree Swallow</b> <i>Tachycineta bicolor</i>                         | <b>13.31%</b>  |
| 293. | <b>Purple Martin</b> <i>Progne subis</i>                               | <b>3.91%</b>   |
| 294. | <b>Northern Rough-winged Swallow</b> <i>Stelgidopteryx serripennis</i> | <b>0.9009%</b> |
| 295. | <b>Barn Swallow</b> <i>Hirundo rustica</i>                             | <b>0.1511%</b> |
| 296. | <b>Cliff Swallow</b> <i>Petrochelidon pyrrhonota</i>                   | <b>0.0006%</b> |
| 297. | <b>Cave Swallow</b> <i>Petrochelidon fulva</i>                         | <b>0.0731%</b> |
| 298. | <b>Red-whiskered Bulbul</b> <i>Pycnonotus jocosus</i>                  | <b>0.0940%</b> |
| 299. | <b>Ruby-crowned Kinglet</b> <i>Corthylio calendula</i>                 | <b>8.85%</b>   |
| 300. | <b>Golden-crowned Kinglet</b> <i>Regulus satrapa</i>                   | <b>0.1920%</b> |
| 301. | <b>White-breasted Nuthatch</b> <i>Sitta carolinensis</i>               | <b>0.4327%</b> |
| 302. | <b>Brown-headed Nuthatch</b> <i>Sitta pusilla</i>                      | <b>1.84%</b>   |
| 303. | <b>Red-breasted Nuthatch</b> <i>Sitta canadensis</i>                   | <b>0.0803%</b> |
| 304. | <b>Brown Creeper</b> <i>Certhia americana</i>                          | <b>0.0197%</b> |
| 305. | <b>Blue-gray Gnatcatcher</b> <i>Poliophtila caerulea</i>               | <b>26.67%</b>  |
| 306. | <b>House Wren</b> <i>Troglodytes aedon</i>                             | <b>6.57%</b>   |
| 307. | <b>Winter Wren</b> <i>Troglodytes hiemalis</i>                         | <b>0.0516%</b> |
| 308. | <b>Sedge Wren</b> <i>Cistothorus stellaris</i>                         | <b>0.7791%</b> |
| 309. | <b>Marsh Wren</b> <i>Cistothorus palustris</i>                         | <b>1.36%</b>   |
| 310. | <b>Carolina Wren</b> <i>Thryothorus ludovicianus</i>                   | <b>23.77%</b>  |

|      |   |         |
|------|---|---------|
| 311. | <b>Bewick's Wren</b> <i>Thryomanes bewickii</i>           | 0.0006% |
| 312. | <b>European Starling</b> <i>Sturnus vulgaris</i>          | 10.45%  |
| 313. | <b>Common Myna</b> <i>Acridotheres tristis</i>            | 0.2100% |
| 314. | <b>Gray Catbird</b> <i>Dumetella carolinensis</i>         | 24.86%  |
| 315. | <b>Brown Thrasher</b> <i>Toxostoma rufum</i>              | 5.23%   |
| 316. | <b>Sage Thrasher</b> <i>Oreoscoptes montanus</i>          | 0.0020% |
| 317. | <b>Bahama Mockingbird</b> <i>Mimus gundlachii</i>         | 0.0366% |
| 318. | <b>Northern Mockingbird</b> <i>Mimus polyglottos</i>      | 37.08%  |
| 319. | <b>Eastern Bluebird</b> <i>Sialia sialis</i>              | 10.26%  |
| 320. | <b>Mountain Bluebird</b> <i>Sialia currucoides</i>        | 0.0131% |
| 321. | <b>Swainson's Thrush</b> <i>Catharus ustulatus</i>        | 0.0017% |
| 322. | <b>Hermit Thrush</b> <i>Catharus guttatus</i>             | 2.03%   |
| 323. | <b>Wood Thrush</b> <i>Hylocichla mustelina</i>            | 0.0104% |
| 324. | <b>American Robin</b> <i>Turdus migratorius</i>           | 18.41%  |
| 325. | <b>Red-legged Thrush</b> <i>Turdus plumbeus</i>           | 0.0003% |
| 326. | <b>Cedar Waxwing</b> <i>Bombycilla cedrorum</i>           | 5.56%   |
| 327. | <b>Tricolored Munia</b> <i>Lonchura malacca</i>           | 0.0220% |
| 328. | <b>House Sparrow</b> <i>Passer domesticus</i>             | 4.34%   |
| 329. | <b>White Wagtail</b> <i>Motacilla alba</i>                | 0.0026% |
| 330. | <b>American Pipit</b> <i>Anthus rubescens</i>             | 0.2956% |
| 331. | <b>Sprague's Pipit</b> <i>Anthus spragueii</i>            | 0.0061% |
| 332. | <b>Evening Grosbeak</b> <i>Coccothraustes vespertinus</i> | 0.0000% |
| 333. | <b>House Finch</b> <i>Haemorhous mexicanus</i>            | 5.83%   |

|      |  |         |
|------|--|---------|
| 334. | <b>Purple Finch</b> <i>Haemorhous purpureus</i>              | 0.2628% |
| 335. | <b>Common Redpoll</b> <i>Acanthis flammea</i>                | 0.0003% |
| 336. | <b>Pine Siskin</b> <i>Spinus pinus</i>                       | 0.4536% |
| 337. | <b>American Goldfinch</b> <i>Spinus tristis</i>              | 7.35%   |
| 338. | <b>Lapland Longspur</b> <i>Calcarius lapponicus</i>          | 0.0035% |
| 339. | <b>Snow Bunting</b> <i>Plectrophenax nivalis</i>             | 0.0041% |
| 340. | <b>Bachman's Sparrow</b> <i>Peucaea aestivalis</i>           | 0.5354% |
| 341. | <b>Grasshopper Sparrow</b> <i>Ammodramus savannarum</i>      | 0.5203% |
| 342. | <b>Chipping Sparrow</b> <i>Spizella passerina</i>            | 7.51%   |
| 343. | <b>Clay-colored Sparrow</b> <i>Spizella pallida</i>          | 0.0943% |
| 344. | <b>Field Sparrow</b> <i>Spizella pusilla</i>                 | 0.3217% |
| 345. | <b>Black-throated Sparrow</b> <i>Amphispiza bilineata</i>    | 0.0000% |
| 346. | <b>Lark Sparrow</b> <i>Chondestes grammacus</i>              | 0.0719% |
| 347. | <b>Lark Bunting</b> <i>Calamospiza melanocorys</i>           | 0.0032% |
| 348. | <b>Fox Sparrow</b> <i>Passerella iliaca</i>                  | 0.0223% |
| 349. | <b>Dark-eyed Junco</b> <i>Junco hyemalis</i>                 | 0.0557% |
| 350. | <b>White-crowned Sparrow</b> <i>Zonotrichia leucophrys</i>   | 0.4063% |
| 351. | <b>Golden-crowned Sparrow</b> <i>Zonotrichia atricapilla</i> | 0.0003% |
| 352. | <b>Harris's Sparrow</b> <i>Zonotrichia querula</i>           | 0.0133% |
| 353. | <b>White-throated Sparrow</b> <i>Zonotrichia albicollis</i>  | 0.6601% |
| 354. | <b>Vesper Sparrow</b> <i>Pooecetes gramineus</i>             | 0.4980% |
| 355. | <b>LeConte's Sparrow</b> <i>Ammospiza leconteii</i>          | 0.0412% |
| 356. | <b>Seaside Sparrow</b> <i>Ammospiza maritima</i>             | 0.0978% |



|      |   |                |
|------|---|----------------|
| 357. | <b>Nelson's Sparrow</b> <i>Ammospiza nelsoni</i>                    | <b>0.1009%</b> |
| 358. | <b>Saltmarsh Sparrow</b> <i>Ammospiza caudacuta</i>                 | <b>0.0742%</b> |
| 359. | <b>Savannah Sparrow</b> <i>Passerculus sandwichensis</i>            | <b>5.11%</b>   |
| 360. | <b>Henslow's Sparrow</b> <i>Centronyx henslowii</i>                 | <b>0.0479%</b> |
| 361. | <b>Song Sparrow</b> <i>Melospiza melodia</i>                        | <b>1.57%</b>   |
| 362. | <b>Lincoln's Sparrow</b> <i>Melospiza lincolnii</i>                 | <b>0.1181%</b> |
| 363. | <b>Swamp Sparrow</b> <i>Melospiza georgiana</i>                     | <b>4.06%</b>   |
| 364. | <b>Green-tailed Towhee</b> <i>Pipilo chlorurus</i>                  | <b>0.0316%</b> |
| 365. | <b>Spotted Towhee</b> <i>Pipilo maculatus</i>                       | <b>0.0006%</b> |
| 366. | <b>Eastern Towhee</b> <i>Pipilo erythrophthalmus</i>                | <b>5.76%</b>   |
| 367. | <b>Western Spindalis</b> <i>Spindalis zena</i>                      | <b>0.0281%</b> |
| 368. | <b>Yellow-breasted Chat</b> <i>Icteria virens</i>                   | <b>0.1047%</b> |
| 369. | <b>Yellow-headed Blackbird</b> <i>Xanthocephalus xanthocephalus</i> | <b>0.1105%</b> |
| 370. | <b>Bobolink</b> <i>Dolichonyx oryzivorus</i>                        | <b>0.0006%</b> |
| 371. | <b>Western Meadowlark</b> <i>Sturnella neglecta</i>                 | <b>0.0064%</b> |
| 372. | <b>Eastern Meadowlark</b> <i>Sturnella magna</i>                    | <b>3.58%</b>   |
| 373. | <b>Orchard Oriole</b> <i>Icterus spurius</i>                        | <b>0.0157%</b> |
| 374. | <b>Hooded Oriole</b> <i>Icterus cucullatus</i>                      | <b>0.0026%</b> |
| 375. | <b>Bullock's Oriole</b> <i>Icterus bullockii</i>                    | <b>0.0386%</b> |
| 376. | <b>Spot-breasted Oriole</b> <i>Icterus pectoralis</i>               | <b>0.2196%</b> |
| 377. | <b>Baltimore Oriole</b> <i>Icterus galbula</i>                      | <b>1.77%</b>   |
| 378. | <b>Scott's Oriole</b> <i>Icterus parisorum</i>                      | <b>0.0125%</b> |

|      |   |                |
|------|---|----------------|
| 379. | <b>Red-winged Blackbird</b> <i>Agelaius phoeniceus</i>      | <b>22.82%</b>  |
| 380. | <b>Tawny-shouldered Blackbird</b> <i>Agelaius humeralis</i> | <b>0.0000%</b> |
| 381. | <b>Shiny Cowbird</b> <i>Molothrus bonariensis</i>           | <b>0.0577%</b> |
| 382. | <b>Bronzed Cowbird</b> <i>Molothrus aeneus</i>              | <b>0.1630%</b> |
| 383. | <b>Brown-headed Cowbird</b> <i>Molothrus ater</i>           | <b>4.68%</b>   |
| 384. | <b>Rusty Blackbird</b> <i>Euphagus carolinus</i>            | <b>0.1096%</b> |
| 385. | <b>Brewer's Blackbird</b> <i>Euphagus cyanocephalus</i>     | <b>0.0255%</b> |
| 386. | <b>Common Grackle</b> <i>Quiscalus quiscula</i>             | <b>21.47%</b>  |
| 387. | <b>Boat-tailed Grackle</b> <i>Quiscalus major</i>           | <b>25.44%</b>  |
| 388. | <b>Great-tailed Grackle</b> <i>Quiscalus mexicanus</i>      | <b>0.0128%</b> |
| 389. | <b>Ovenbird</b> <i>Seiurus aurocapilla</i>                  | <b>1.33%</b>   |
| 390. | <b>Worm-eating Warbler</b> <i>Helmitheros vermivorum</i>    | <b>0.0745%</b> |
| 391. | <b>Louisiana Waterthrush</b> <i>Parkesia motacilla</i>      | <b>0.1270%</b> |
| 392. | <b>Northern Waterthrush</b> <i>Parkesia noveboracensis</i>  | <b>0.6195%</b> |
| 393. | <b>Golden-winged Warbler</b> <i>Vermivora chrysoptera</i>   | <b>0.0067%</b> |
| 394. | <b>Blue-winged Warbler</b> <i>Vermivora cyanoptera</i>      | <b>0.0107%</b> |
| 395. | <b>Black-and-white Warbler</b> <i>Mniotilta varia</i>       | <b>7.05%</b>   |
| 396. | <b>Prothonotary Warbler</b> <i>Protonotaria citrea</i>      | <b>0.0044%</b> |
| 397. | <b>Swainson's Warbler</b> <i>Limnothlypis swainsonii</i>    | <b>0.0162%</b> |
| 398. | <b>Tennessee Warbler</b> <i>Leiothlypis peregrina</i>       | <b>0.0049%</b> |
| 399. | <b>Orange-crowned Warbler</b> <i>Leiothlypis celata</i>     | <b>3.19%</b>   |
| 400. | <b>Nashville Warbler</b> <i>Leiothlypis ruficapilla</i>     | <b>0.0696%</b> |
| 401. | <b>MacGillivray's Warbler</b> <i>Geothlypis tolmiei</i>     | <b>0.0119%</b> |

|  |         |
|--|---------|
| 402. <b>Common Yellowthroat</b> <i>Geothlypis trichas</i>              | 10.07%  |
| 403. <b>Hooded Warbler</b> <i>Setophaga citrina</i>                    | 0.0131% |
| 404. <b>American Redstart</b> <i>Setophaga ruticilla</i>               | 0.8936% |
| 405. <b>Kirtland's Warbler</b> <i>Setophaga kirtlandii</i>             | 0.0058% |
| 406. <b>Cape May Warbler</b> <i>Setophaga tigrina</i>                  | 0.1128% |
| 407. <b>Northern Parula</b> <i>Setophaga americana</i>                 | 8.04%   |
| 408. <b>Magnolia Warbler</b> <i>Setophaga magnolia</i>                 | 0.2106% |
| 409. <b>Bay-breasted Warbler</b> <i>Setophaga castanea</i>             | 0.0055% |
| 410. <b>Blackburnian Warbler</b> <i>Setophaga fusca</i>                | 0.0012% |
| 411. <b>Yellow Warbler</b> <i>Setophaga petechia</i>                   | 0.1653% |
| 412. <b>Chestnut-sided Warbler</b> <i>Setophaga pensylvanica</i>       | 0.0012% |
| 413. <b>Blackpoll Warbler</b> <i>Setophaga striata</i>                 | 0.0009% |
| 414. <b>Black-throated Blue Warbler</b> <i>Setophaga caeruleascens</i> | 0.2706% |
| 415. <b>Palm Warbler</b> <i>Setophaga palmarum</i>                     | 37.06%  |
| 416. <b>Pine Warbler</b> <i>Setophaga pinus</i>                        | 12.62%  |
| 417. <b>Yellow-rumped Warbler</b> <i>Setophaga coronata</i>            | 43.54%  |
| 418. <b>Yellow-throated Warbler</b> <i>Setophaga dominica</i>          | 6.04%   |
| 419. <b>Prairie Warbler</b> <i>Setophaga discolor</i>                  | 3.66%   |
| 420. <b>Black-throated Gray Warbler</b> <i>Setophaga nigrescens</i>    | 0.0241% |
| 421. <b>Townsend's Warbler</b> <i>Setophaga townsendi</i>              | 0.0009% |
| 422. <b>Black-throated Green Warbler</b> <i>Setophaga virens</i>       | 0.5543% |
| 423. <b>Wilson's Warbler</b> <i>Cardellina pusilla</i>                 | 0.0650% |

|                            |   |                          |
|----------------------------|---|--------------------------|
| 424.                       | <b>Summer Tanager</b> <i>Piranga rubra</i>                    | <b>0.5836%</b>           |
| 425.                       | <b>Scarlet Tanager</b> <i>Piranga olivacea</i>                | <b>0.0017%</b>           |
| 426.                       | <b>Western Tanager</b> <i>Piranga ludoviciana</i>             | <b>0.1717%</b>           |
| 427.                       | <b>Northern Cardinal</b> <i>Cardinalis cardinalis</i>         | <b>47.58%</b>            |
| 428.                       | <b>Rose-breasted Grosbeak</b> <i>Pheucticus ludovicianus</i>  | <b>0.0618%</b>           |
| 429.                       | <b>Black-headed Grosbeak</b> <i>Pheucticus melanocephalus</i> | <b>0.0110%</b>           |
| 430.                       | <b>Blue Grosbeak</b> <i>Passerina caerulea</i>                | <b>0.1204%</b>           |
| 431.                       | <b>Lazuli Bunting</b> <i>Passerina amoena</i>                 | <b>0.0122%</b>           |
| 432.                       | <b>Indigo Bunting</b> <i>Passerina cyanea</i>                 | <b>1.25%</b>             |
| 433.                       | <b>Painted Bunting</b> <i>Passerina ciris</i>                 | <b>6.42%</b>             |
| 434.                       | <b>Dickcissel</b> <i>Spiza americana</i>                      | <b>0.0464%</b>           |
| 435.                       | <b>Red-legged Honeycreeper</b> <i>Cyanerpes cyaneus</i>       | <b>0.0000%</b>           |
| 436.                       | <b>Bananaquit</b> <i>Coereba flaveola</i>                     | <b>0.0238%</b>           |
| 437.                       | <b>Yellow-faced Grassquit</b> <i>Tiaris olivaceus</i>         | <b>0.0023%</b>           |
| 438.                       | <b>Black-faced Grassquit</b> <i>Melanospiza bicolor</i>       | <b>0.0264%</b>           |
| <b>EXOTIC: PROVISIONAL</b> |   |                          |
| 439.                       | <b>Indian Peafowl</b> <i>Pavo cristatus</i>                   | <b>0.3362%</b> frequency |
| 440.                       | <b>Red Junglefowl</b> <i>Gallus gallus</i>                    | <b>0.0922%</b>           |
| 441.                       | <b>Great White Pelican</b> <i>Pelecanus onocrotalus</i>       | <b>0.0220%</b>           |
| 442.                       | <b>Rose-ringed Parakeet</b> <i>Psittacula krameri</i>         | <b>0.0304%</b>           |
| 443.                       | <b>Yellow-chevroned Parakeet</b> <i>Brotogeris chiriri</i>    | <b>0.1856%</b>           |
| 444.                       | <b>Red-crowned Parrot</b> <i>Amazona viridigenalis</i>        | <b>0.0499%</b>           |

|   |                |
|---|----------------|
| 445. <b>Orange-winged Parrot</b> <i>Amazona amazonica</i>           | <b>0.0832%</b> |
| 446. <b>Blue-and-yellow Macaw</b> <i>Ara ararauna</i>               | <b>0.0629%</b> |
| 447. <b>Chestnut-fronted Macaw</b> <i>Ara severus</i>               | <b>0.0490%</b> |
| 448. <b>Blue-crowned Parakeet</b> <i>Thectocercus acuticaudatus</i> | <b>0.2938%</b> |
| 449. <b>Red-masked Parakeet</b> <i>Psittacara erythrogenys</i>      | <b>0.1091%</b> |
| 450. <b>White-eyed Parakeet</b> <i>Psittacara leucophthalmus</i>    | <b>0.0310%</b> |
| 451. <b>House Crow</b> <i>Corvus splendens</i>                      | <b>0.0012%</b> |
| 452. <b>Common Hill Myna</b> <i>Gracula religiosa</i>               | <b>0.0418%</b> |
| 453. <b>Scaly-breasted Munia</b> <i>Lonchura punctulata</i>         | <b>0.2169%</b> |

**EXOTIC: ESCAPEE**

|  |                          |
|--|--------------------------|
| <b>Emu</b> <i>Dromaius novaehollandiae</i>                   | <b>0.0000%</b> frequency |
| <b>Plumed Whistling-Duck</b> <i>Dendrocygna eytoni</i>       | <b>0.0006%</b>           |
| <b>White-faced Whistling-Duck</b> <i>Dendrocygna viduata</i> | <b>0.0136%</b>           |
| <b>Bar-headed Goose</b> <i>Anser indicus</i>                 | <b>0.0020%</b>           |
| <b>Graylag Goose</b> <i>Anser anser</i>                      | <b>0.1555%</b>           |
| <b>Swan Goose</b> <i>Anser cygnoides</i>                     | <b>0.1328%</b>           |
| <b>Barnacle Goose</b> <i>Branta leucopsis</i>                | <b>0.0006%</b>           |
| <b>Hawaiian Goose</b> <i>Branta sandvicensis</i>             | <b>0.0003%</b>           |
| <b>Mute Swan</b> <i>Cygnus olor</i>                          | <b>0.2758%</b>           |
| <b>Black Swan</b> <i>Cygnus atratus</i>                      | <b>0.1166%</b>           |
| <b>Black-necked Swan</b> <i>Cygnus melancoryphus</i>         | <b>0.0421%</b>           |

|  |                |
|--|----------------|
| <b>Trumpeter Swan</b> <i>Cygnus buccinator</i>               | <b>0.0206%</b> |
| <b>Whooper Swan</b> <i>Cygnus cygnus</i>                     | <b>0.0148%</b> |
| <b>Coscoroba Swan</b> <i>Coscoroba coscoroba</i>             | <b>0.0102%</b> |
| <b>Radjah Shelduck</b> <i>Radjah radjah</i>                  | <b>0.0003%</b> |
| <b>Ruddy Shelduck</b> <i>Tadorna ferruginea</i>              | <b>0.0064%</b> |
| <b>Common Shelduck</b> <i>Tadorna tadorna</i>                | <b>0.0133%</b> |
| <b>Ringed Teal</b> <i>Callonetta leucophrys</i>              | <b>0.0006%</b> |
| <b>Mandarin Duck</b> <i>Aix galericulata</i>                 | <b>0.0168%</b> |
| <b>Red-crested Pochard</b> <i>Netta rufina</i>               | <b>0.0014%</b> |
| <b>Rosy-billed Pochard</b> <i>Netta peposaca</i>             | <b>0.0003%</b> |
| <b>Common Pochard</b> <i>Aythya ferina</i>                   | <b>0.0000%</b> |
| <b>Helmeted Guineafowl</b> <i>Numida meleagris</i>           | <b>0.0334%</b> |
| <b>Ring-necked Pheasant</b> <i>Phasianus colchicus</i>       | <b>0.0046%</b> |
| <b>Silver Pheasant</b> <i>Lophura nycthemera</i>             | <b>0.0000%</b> |
| <b>Green Peafowl</b> <i>Pavo muticus</i>                     | <b>0.0009%</b> |
| <b>Red-legged Partridge</b> <i>Alectoris rufa</i>            | <b>0.0046%</b> |
| <b>African Collared-Dove</b> <i>Streptopelia roseogrisea</i> | <b>0.0017%</b> |
| <b>Ring-necked Dove</b> <i>Streptopelia capicola</i>         | <b>0.0000%</b> |
| <b>Giant Wood-Rail</b> <i>Aramides ypecaha</i>               | <b>0.0017%</b> |
| <b>Black-necked Stork</b> <i>Ephippiorhynchus asiaticus</i>  | <b>0.0003%</b> |
| <b>African Sacred Ibis</b> <i>Threskiornis aethiopicus</i>   | <b>0.0003%</b> |
| <b>Great Black Hawk</b> <i>Buteogallus urubitinga</i>        | <b>0.0017%</b> |

|  |                |
|--|----------------|
| <b>Harris's Hawk</b> <i>Parabuteo unicinctus</i>         | <b>0.0003%</b> |
| <b>Cockatiel</b> <i>Nymphicus hollandicus</i>            | <b>0.0029%</b> |
| <b>Chattering Lory</b> <i>Lorius garrulus</i>            | <b>0.0000%</b> |
| <b>Rosy-faced Lovebird</b> <i>Agapornis roseicollis</i>  | <b>0.0009%</b> |
| <b>Fischer's Lovebird</b> <i>Agapornis fischeri</i>      | <b>0.0000%</b> |
| <b>Gray Parrot</b> <i>Psittacus erithacus</i>            | <b>0.0000%</b> |
| <b>Scaly-headed Parrot</b> <i>Pionus maximiliani</i>     | <b>0.0012%</b> |
| <b>Lilac-crowned Parrot</b> <i>Amazona finschi</i>       | <b>0.0046%</b> |
| <b>Red-lored Parrot</b> <i>Amazona autumnalis</i>        | <b>0.0003%</b> |
| <b>Yellow-naped Parrot</b> <i>Amazona auropalliata</i>   | <b>0.0006%</b> |
| <b>Yellow-headed Parrot</b> <i>Amazona oratrix</i>       | <b>0.0009%</b> |
| <b>Yellow-crowned Parrot</b> <i>Amazona ochrocephala</i> | <b>0.0078%</b> |
| <b>Turquoise-fronted Parrot</b> <i>Amazona aestiva</i>   | <b>0.0058%</b> |
| <b>White-fronted Parrot</b> <i>Amazona albifrons</i>     | <b>0.0017%</b> |
| <b>Mealy Parrot</b> <i>Amazona farinosa</i>              | <b>0.0006%</b> |
| <b>Hyacinth Macaw</b> <i>Anodorhynchus hyacinthinus</i>  | <b>0.0003%</b> |
| <b>Dusky-headed Parakeet</b> <i>Aratinga weddellii</i>   | <b>0.0000%</b> |
| <b>Sun Parakeet</b> <i>Aratinga solstitialis</i>         | <b>0.0003%</b> |
| <b>Military Macaw</b> <i>Ara militaris</i>               | <b>0.0000%</b> |
| <b>Red-and-green Macaw</b> <i>Ara chloropterus</i>       | <b>0.0003%</b> |
| <b>Golden Parakeet</b> <i>Guaruba guarouba</i>           | <b>0.0000%</b> |
| <b>Green Parakeet</b> <i>Psittacara holochlorus</i>      | <b>0.0052%</b> |

|   |                |
|---|----------------|
| <b>Crimson-fronted Parakeet</b> <i>Psittacara finschi</i> | <b>0.0078%</b> |
| <b>Black-throated Magpie-Jay</b> <i>Calocitta colliei</i> | <b>0.0000%</b> |
| <b>Tropical Mockingbird</b> <i>Mimus gilvus</i>           | <b>0.0012%</b> |
| <b>Northern Red Bishop</b> <i>Euplectes franciscanus</i>  | <b>0.0003%</b> |
| <b>Zebra Finch</b> <i>Taeniopygia guttata</i>             | <b>0.0012%</b> |
| <b>Bronze Mannikin</b> <i>Spermestes cucullata</i>        | <b>0.0000%</b> |
| <b>Yellow-fronted Canary</b> <i>Crithagra mozambica</i>   | <b>0.0000%</b> |
| <b>Saffron Finch</b> <i>Sicalis flaveola</i>              | <b>0.0003%</b> |
| <b>Scarlet Ibis</b> <i>Eudocimus ruber</i>                | <b>0.0000%</b> |

**HYBRIDS**

|   |                          |
|---|--------------------------|
| Graylag x Swan Goose (hybrid) <i>Anser anser x cygnoides</i>                                    | <b>0.0102%</b> frequency |
| Swan Goose x Canada Goose (hybrid) <i>Anser cygnoides x Branta canadensis</i>                   | <b>0.0014%</b>           |
| Graylag x Canada Goose (hybrid) <i>Anser anser x Branta canadensis</i>                          | <b>0.0009%</b>           |
| Domestic goose sp. x Canada Goose (hybrid) <i>Anser sp. (Domestic type) x Branta canadensis</i> | <b>0.0006%</b>           |
| Blue-winged x Cinnamon Teal (hybrid) <i>Spatula discors x cyanoptera</i>                        | <b>0.0145%</b>           |
| Blue-winged Teal x Northern Shoveler (hybrid) <i>Spatula discors x clypeata</i>                 | <b>0.0003%</b>           |
| Eurasian x American Wigeon (hybrid) <i>Mareca penelope x americana</i>                          | <b>0.0009%</b>           |
| Muscovy Duck x Mallard (hybrid) <i>Cairina moschata x Anas platyrhynchos</i>                    | <b>0.0415%</b>           |
| Northern Shoveler x Mallard (hybrid) <i>Spatula clypeata x Anas platyrhynchos</i>               | <b>0.0003%</b>           |



|   |         |
|---|---------|
| Mallard x Eastern Spot-billed Duck (hybrid)<br><i>Anas platyrhynchos x zonorhyncha</i>    | 0.0009% |
| Mallard x American Black Duck (hybrid) <i>Anas platyrhynchos x rubripes</i>               | 0.0003% |
| Mallard x Mottled Duck (hybrid) <i>Anas platyrhynchos x fulvigula</i>                     | 1.76%   |
| Redhead x Lesser Scaup (hybrid) <i>Aythya americana x affinis</i>                         | 0.0009% |
| Rufous x Allen's Hummingbird (hybrid)<br><i>Selasphorus rufus x sasin</i>                 | 0.0012% |
| Herring x Great Black-backed Gull (hybrid) <i>Larus argentatus x marinus</i>              | 0.0006% |
| Herring x Glaucous Gull (hybrid) <i>Larus argentatus x hyperboreus</i>                    | 0.0038% |
| Herring x Lesser Black-backed Gull (hybrid) <i>Larus argentatus x fuscus</i>              | 0.0026% |
| Double-crested x Neotropic Cormorant (hybrid)<br><i>Nannopterum auritum x brasilianum</i> | 0.0049% |
| Little Blue x Tricolored Heron (hybrid) <i>Egretta caerulea x tricolor</i>                | 0.0003% |
| Tricolored Heron x Snowy Egret (hybrid) <i>Egretta tricolor x thula</i>                   | 0.0006% |
| Great Egret x Great Blue Heron (hybrid) <i>Ardea alba x herodias</i>                      | 0.0049% |
| White x Scarlet Ibis (hybrid) <i>Eudocimus albus x ruber</i>                              | 0.0012% |
| Blue-and-yellow x Scarlet Macaw (hybrid)<br><i>Ara ararauna x macao</i>                   | 0.0000% |
| Tropical x Northern Mockingbird (hybrid)<br><i>Mimus gilvus x polyglottos</i>             | 0.0003% |
| Indigo x Painted Bunting (hybrid) <i>Passerina cyanea x ciris</i>                         | 0.0023% |