



Target Species

Based on 313,100 complete checklists

465 species observed in Florida, US (region/US-FL) in Mar that you need for your **Florida Life List**

| | |
|--|------------------------|
| 1. Black-bellied Whistling-Duck <i>Dendrocygna autumnalis</i> | 3.99% frequency |
| 2. Fulvous Whistling-Duck <i>Dendrocygna bicolor</i> | 0.4650% |
| 3. Snow Goose <i>Anser caerulescens</i> | 0.0588% |
| 4. Ross's Goose <i>Anser rossii</i> | 0.0077% |
| 5. Greater White-fronted Goose <i>Anser albifrons</i> | 0.0224% |
| 6. Brant <i>Branta bernicla</i> | 0.0061% |
| 7. Canada Goose <i>Branta canadensis</i> | 2.44% |
| 8. Tundra Swan <i>Cygnus columbianus</i> | 0.0010% |
| 9. Egyptian Goose <i>Alopochen aegyptiaca</i> | 1.61% |
| 10. Muscovy Duck <i>Cairina moschata</i> | 4.50% |
| 11. Wood Duck <i>Aix sponsa</i> | 4.66% |
| 12. Blue-winged Teal <i>Spatula discors</i> | 10.11% |
| 13. Cinnamon Teal <i>Spatula cyanoptera</i> | 0.0786% |
| 14. Northern Shoveler <i>Spatula clypeata</i> | 2.09% |
| 15. Gadwall <i>Mareca strepera</i> | 0.2913% |
| 16. Eurasian Wigeon <i>Mareca penelope</i> | 0.0211% |
| 17. American Wigeon <i>Mareca americana</i> | 0.7656% |

| | | |
|-----|--|---------|
| 18. | Mallard <i>Anas platyrhynchos</i> | 3.53% |
| 19. | American Black Duck <i>Anas rubripes</i> | 0.0109% |
| 20. | Mottled Duck <i>Anas fulvigula</i> | 11.36% |
| 21. | White-cheeked Pintail <i>Anas bahamensis</i> | 0.0182% |
| 22. | Northern Pintail <i>Anas acuta</i> | 0.1507% |
| 23. | Green-winged Teal <i>Anas crecca</i> | 0.8905% |
| 24. | Canvasback <i>Aythya valisineria</i> | 0.0824% |
| 25. | Redhead <i>Aythya americana</i> | 0.7122% |
| 26. | Ring-necked Duck <i>Aythya collaris</i> | 2.12% |
| 27. | Greater Scaup <i>Aythya marila</i> | 0.2830% |
| 28. | Lesser Scaup <i>Aythya affinis</i> | 3.01% |
| 29. | Common Eider <i>Somateria mollissima</i> | 0.0131% |
| 30. | Harlequin Duck <i>Histrionicus histrionicus</i> | 0.0089% |
| 31. | Surf Scoter <i>Melanitta perspicillata</i> | 0.0374% |
| 32. | White-winged Scoter <i>Melanitta deglandi</i> | 0.0239% |
| 33. | Black Scoter <i>Melanitta americana</i> | 0.1453% |
| 34. | Long-tailed Duck <i>Clangula hyemalis</i> | 0.0236% |
| 35. | Bufflehead <i>Bucephala albeola</i> | 1.40% |
| 36. | Common Goldeneye <i>Bucephala clangula</i> | 0.0431% |
| 37. | Hooded Merganser <i>Lophodytes cucullatus</i> | 1.33% |
| 38. | Common Merganser <i>Mergus merganser</i> | 0.0010% |
| 39. | Red-breasted Merganser <i>Mergus serrator</i> | 3.69% |
| 40. | Masked Duck <i>Nomonyx dominicus</i> | 0.0006% |

| | |
|---|----------------|
| 41. Ruddy Duck <i>Oxyura jamaicensis</i> | 0.4973% |
| 42. Northern Bobwhite <i>Colinus virginianus</i> | 1.77% |
| 43. Wild Turkey <i>Meleagris gallopavo</i> | 2.99% |
| 44. American Flamingo <i>Phoenicopterus ruber</i> | 0.1616% |
| 45. Least Grebe <i>Tachybaptus dominicus</i> | 0.0444% |
| 46. Pied-billed Grebe <i>Podilymbus podiceps</i> | 12.65% |
| 47. Horned Grebe <i>Podiceps auritus</i> | 0.5634% |
| 48. Red-necked Grebe <i>Podiceps grisegena</i> | 0.0112% |
| 49. Eared Grebe <i>Podiceps nigricollis</i> | 0.0086% |
| 50. Western Grebe <i>Aechmophorus occidentalis</i> | 0.0003% |
| 51. Rock Pigeon <i>Columba livia</i> | 6.77% |
| 52. White-crowned Pigeon <i>Patagioenas leucocephala</i> | 0.5529% |
| 53. Eurasian Collared-Dove <i>Streptopelia decaocto</i> | 11.42% |
| 54. Inca Dove <i>Columbina inca</i> | 0.0029% |
| 55. Common Ground Dove <i>Columbina passerina</i> | 9.38% |
| 56. Key West Quail-Dove <i>Geotrygon chrysia</i> | 0.0077% |
| 57. White-winged Dove <i>Zenaida asiatica</i> | 4.14% |
| 58. Zenaida Dove <i>Zenaida aurita</i> | 0.0249% |
| 59. Mourning Dove <i>Zenaida macroura</i> | 43.82% |
| 60. Smooth-billed Ani <i>Crotophaga ani</i> | 0.0968% |
| 61. Groove-billed Ani <i>Crotophaga sulcirostris</i> | 0.0505% |
| 62. Yellow-billed Cuckoo <i>Coccyzus americanus</i> | 0.0639% |
| 63. Mangrove Cuckoo <i>Coccyzus minor</i> | 0.0974% |

| | | |
|-----|---|----------------|
| 64. | Black-billed Cuckoo <i>Coccyzus erythrophthalmus</i> | 0.0003% |
| 65. | Lesser Nighthawk <i>Chordeiles acutipennis</i> | 0.0128% |
| 66. | Common Nighthawk <i>Chordeiles minor</i> | 0.0115% |
| 67. | Chuck-will's-widow <i>Antrostomus carolinensis</i> | 1.42% |
| 68. | Eastern Whip-poor-will <i>Antrostomus vociferus</i> | 0.3245% |
| 69. | Chimney Swift <i>Chaetura pelagica</i> | 0.5573% |
| 70. | Vaux's Swift <i>Chaetura vauxi</i> | 0.0093% |
| 71. | Ruby-throated Hummingbird <i>Archilochus colubris</i> | 4.27% |
| 72. | Black-chinned Hummingbird <i>Archilochus alexandri</i> | 0.0313% |
| 73. | Calliope Hummingbird <i>Selasphorus calliope</i> | 0.0160% |
| 74. | Rufous Hummingbird <i>Selasphorus rufus</i> | 0.0767% |
| 75. | Allen's Hummingbird <i>Selasphorus sasin</i> | 0.0003% |
| 76. | Broad-billed Hummingbird <i>Cynanthus latirostris</i> | 0.0006% |
| 77. | Buff-bellied Hummingbird <i>Amazilia yucatanensis</i> | 0.0476% |
| 78. | King Rail <i>Rallus elegans</i> | 0.5889% |
| 79. | Clapper Rail <i>Rallus crepitans</i> | 1.43% |
| 80. | Virginia Rail <i>Rallus limicola</i> | 0.3053% |
| 81. | Sora <i>Porzana carolina</i> | 2.47% |
| 82. | Common Gallinule <i>Gallinula galeata</i> | 19.89% |
| 83. | American Coot <i>Fulica americana</i> | 12.58% |
| 84. | Purple Gallinule <i>Porphyrio martinica</i> | 4.43% |
| 85. | Gray-headed Swamphen <i>Porphyrio poliocephalus</i> | 1.67% |

| | |
|---|---------|
| 86. Yellow Rail <i>Coturnicops noveboracensis</i> | 0.0016% |
| 87. Black Rail <i>Laterallus jamaicensis</i> | 0.0326% |
| 88. Limpkin <i>Aramus guarauna</i> | 9.88% |
| 89. Sandhill Crane <i>Antigone canadensis</i> | 10.36% |
| 90. Whooping Crane <i>Grus americana</i> | 0.1469% |
| 91. Black-necked Stilt <i>Himantopus mexicanus</i> | 3.91% |
| 92. American Avocet <i>Recurvirostra americana</i> | 0.7125% |
| 93. American Oystercatcher <i>Haematopus palliatus</i> | 1.39% |
| 94. Black-bellied Plover <i>Pluvialis squatarola</i> | 4.61% |
| 95. American Golden-Plover <i>Pluvialis dominica</i> | 0.0351% |
| 96. Pacific Golden-Plover <i>Pluvialis fulva</i> | 0.0026% |
| 97. Killdeer <i>Charadrius vociferus</i> | 8.53% |
| 98. Semipalmated Plover <i>Charadrius semipalmatus</i> | 2.63% |
| 99. Piping Plover <i>Charadrius melodus</i> | 0.8116% |
| 100. Wilson's Plover <i>Anarhynchus wilsonia</i> | 1.14% |
| 101. Snowy Plover <i>Anarhynchus nivosus</i> | 0.5509% |
| 102. Upland Sandpiper <i>Bartramia longicauda</i> | 0.0077% |
| 103. Whimbrel <i>Numenius phaeopus</i> | 0.2118% |
| 104. Long-billed Curlew <i>Numenius americanus</i> | 0.0620% |
| 105. Bar-tailed Godwit <i>Limosa lapponica</i> | 0.0013% |
| 106. Black-tailed Godwit <i>Limosa limosa</i> | 0.0003% |
| 107. Marbled Godwit <i>Limosa fedoa</i> | 1.01% |
| 108. Short-billed Dowitcher <i>Limnodromus griseus</i> | 2.22% |

| | | |
|------|--|---------|
| 109. | Long-billed Dowitcher <i>Limnodromus scolopaceus</i> | 1.16% |
| 110. | American Woodcock <i>Scolopax minor</i> | 0.0278% |
| 111. | Wilson's Snipe <i>Gallinago delicata</i> | 2.36% |
| 112. | Wilson's Phalarope <i>Phalaropus tricolor</i> | 0.0022% |
| 113. | Red Phalarope <i>Phalaropus fulicarius</i> | 0.0016% |
| 114. | Spotted Sandpiper <i>Actitis macularius</i> | 2.40% |
| 115. | Solitary Sandpiper <i>Tringa solitaria</i> | 0.6426% |
| 116. | Lesser Yellowlegs <i>Tringa flavipes</i> | 4.05% |
| 117. | Willet <i>Tringa semipalmata</i> | 7.96% |
| 118. | Greater Yellowlegs <i>Tringa melanoleuca</i> | 5.05% |
| 119. | Ruddy Turnstone <i>Arenaria interpres</i> | 5.40% |
| 120. | Red Knot <i>Calidris canutus</i> | 0.7285% |
| 121. | Ruff <i>Calidris pugnax</i> | 0.0038% |
| 122. | Curlew Sandpiper <i>Calidris ferruginea</i> | 0.0000% |
| 123. | Stilt Sandpiper <i>Calidris himantopus</i> | 0.4436% |
| 124. | Buff-breasted Sandpiper <i>Calidris subruficollis</i> | 0.0006% |
| 125. | Sanderling <i>Calidris alba</i> | 5.69% |
| 126. | Dunlin <i>Calidris alpina</i> | 3.04% |
| 127. | Purple Sandpiper <i>Calidris maritima</i> | 0.0543% |
| 128. | Baird's Sandpiper <i>Calidris bairdii</i> | 0.0000% |
| 129. | White-rumped Sandpiper <i>Calidris fuscicollis</i> | 0.0003% |
| 130. | Least Sandpiper <i>Calidris minutilla</i> | 4.19% |
| 131. | Pectoral Sandpiper <i>Calidris melanotos</i> | 0.1578% |

| | |
|--|---------|
| 132. Western Sandpiper <i>Calidris mauri</i> | 1.38% |
| 133. Semipalmated Sandpiper <i>Calidris pusilla</i> | 0.0364% |
| 134. Parasitic Jaeger <i>Stercorarius parasiticus</i> | 0.0326% |
| 135. Pomarine Jaeger <i>Stercorarius pomarinus</i> | 0.0169% |
| 136. Long-billed Murrelet <i>Brachyramphus perdix</i> | 0.0010% |
| 137. Razorbill <i>Alca torda</i> | 0.0013% |
| 138. Little Gull <i>Hydrocoloeus minutus</i> | 0.0003% |
| 139. Black-legged Kittiwake <i>Rissa tridactyla</i> | 0.0128% |
| 140. Bonaparte's Gull <i>Chroicocephalus philadelphia</i> | 1.59% |
| 141. Black-headed Gull <i>Chroicocephalus ridibundus</i> | 0.0051% |
| 142. Laughing Gull <i>Leucophaeus atricilla</i> | 19.49% |
| 143. Franklin's Gull <i>Leucophaeus pipixcan</i> | 0.0086% |
| 144. Heermann's Gull <i>Larus heermanni</i> | 0.0054% |
| 145. Ring-billed Gull <i>Larus delawarensis</i> | 11.68% |
| 146. Herring Gull <i>Larus argentatus</i> | 3.97% |
| 147. Yellow-legged Gull <i>Larus michahellis</i> | 0.0003% |
| 148. Great Black-backed Gull <i>Larus marinus</i> | 0.5698% |
| 149. Glaucous Gull <i>Larus hyperboreus</i> | 0.0559% |
| 150. Lesser Black-backed Gull <i>Larus fuscus</i> | 1.45% |
| 151. California Gull <i>Larus californicus</i> | 0.0035% |
| 152. Slaty-backed Gull <i>Larus schistisagus</i> | 0.0000% |
| 153. Iceland Gull <i>Larus glaucoides</i> | 0.0409% |
| 154. Black Skimmer <i>Rynchops niger</i> | 3.31% |

| | | |
|------|---|---------|
| 155. | Brown Noddy <i>Anous stolidus</i> | 0.1070% |
| 156. | Black Noddy <i>Anous minutus</i> | 0.0010% |
| 157. | Sooty Tern <i>Onychoprion fuscatus</i> | 0.1153% |
| 158. | Bridled Tern <i>Onychoprion anaethetus</i> | 0.0045% |
| 159. | Least Tern <i>Sternula antillarum</i> | 0.3635% |
| 160. | Gull-billed Tern <i>Gelochelidon nilotica</i> | 0.1140% |
| 161. | Caspian Tern <i>Hydroprogne caspia</i> | 2.94% |
| 162. | Black Tern <i>Chlidonias niger</i> | 0.0067% |
| 163. | Forster's Tern <i>Sterna forsteri</i> | 6.09% |
| 164. | Common Tern <i>Sterna hirundo</i> | 0.1798% |
| 165. | Roseate Tern <i>Sterna dougallii</i> | 0.0058% |
| 166. | Sandwich Tern <i>Thalasseus sandvicensis</i> | 2.00% |
| 167. | Elegant Tern <i>Thalasseus elegans</i> | 0.0026% |
| 168. | Royal Tern <i>Thalasseus maximus</i> | 8.75% |
| 169. | Red-billed Tropicbird <i>Phaethon aethereus</i> | 0.0016% |
| 170. | Red-throated Loon <i>Gavia stellata</i> | 0.0160% |
| 171. | Pacific Loon <i>Gavia pacifica</i> | 0.0070% |
| 172. | Common Loon <i>Gavia immer</i> | 2.20% |
| 173. | Wilson's Storm-Petrel <i>Oceanites oceanicus</i> | 0.0013% |
| 174. | Black-capped Petrel <i>Pterodroma hasitata</i> | 0.0003% |
| 175. | Cory's Shearwater <i>Calonectris diomedea</i> | 0.0016% |
| 176. | Great Shearwater <i>Ardenna gravis</i> | 0.0003% |
| 177. | Sooty Shearwater <i>Ardenna grisea</i> | 0.0006% |

| | | |
|------|--|---------|
| 178. | Audubon's Shearwater <i>Puffinus lherminieri</i> | 0.0026% |
| 179. | Wood Stork <i>Mycteria americana</i> | 12.13% |
| 180. | Magnificent Frigatebird <i>Fregata magnificens</i> | 1.79% |
| 181. | Northern Gannet <i>Morus bassanus</i> | 0.9959% |
| 182. | Brown Booby <i>Sula leucogaster</i> | 0.1226% |
| 183. | Masked Booby <i>Sula dactylatra</i> | 0.0591% |
| 184. | Anhinga <i>Anhinga anhinga</i> | 25.32% |
| 185. | Great Cormorant <i>Phalacrocorax carbo</i> | 0.0121% |
| 186. | Double-crested Cormorant <i>Nannopterum auritum</i> | 31.30% |
| 187. | Neotropic Cormorant <i>Nannopterum brasilianum</i> | 0.0693% |
| 188. | American White Pelican <i>Pelecanus erythrorhynchos</i> | 5.71% |
| 189. | Brown Pelican <i>Pelecanus occidentalis</i> | 17.74% |
| 190. | American Bittern <i>Botaurus lentiginosus</i> | 1.52% |
| 191. | Least Bittern <i>Ixobrychus exilis</i> | 1.15% |
| 192. | Yellow-crowned Night Heron <i>Nyctanassa violacea</i> | 2.74% |
| 193. | Black-crowned Night Heron <i>Nycticorax nycticorax</i> | 5.09% |
| 194. | Little Blue Heron <i>Egretta caerulea</i> | 25.31% |
| 195. | Tricolored Heron <i>Egretta tricolor</i> | 20.53% |
| 196. | Reddish Egret <i>Egretta rufescens</i> | 2.91% |
| 197. | Snowy Egret <i>Egretta thula</i> | 20.57% |
| 198. | Green Heron <i>Butorides virescens</i> | 10.90% |
| 199. | Western Cattle Egret <i>Bubulcus ibis</i> | 13.02% |

| | | |
|------|---|----------------|
| 200. | Great Egret <i>Ardea alba</i> | 33.47% |
| 201. | Great Blue Heron <i>Ardea herodias</i> | 30.48% |
| 202. | White Ibis <i>Eudocimus albus</i> | 34.48% |
| 203. | Glossy Ibis <i>Plegadis falcinellus</i> | 12.21% |
| 204. | White-faced Ibis <i>Plegadis chihi</i> | 0.0920% |
| 205. | Roseate Spoonbill <i>Platalea ajaja</i> | 6.41% |
| 206. | Black Vulture <i>Coragyps atratus</i> | 25.97% |
| 207. | Turkey Vulture <i>Cathartes aura</i> | 40.24% |
| 208. | Osprey <i>Pandion haliaetus</i> | 34.15% |
| 209. | White-tailed Kite <i>Elanus leucurus</i> | 0.1204% |
| 210. | Swallow-tailed Kite <i>Elanoides forficatus</i> | 9.19% |
| 211. | Golden Eagle <i>Aquila chrysaetos</i> | 0.0003% |
| 212. | Snail Kite <i>Rostrhamus sociabilis</i> | 1.41% |
| 213. | Mississippi Kite <i>Ictinia mississippiensis</i> | 0.0112% |
| 214. | Northern Harrier <i>Circus hudsonius</i> | 4.93% |
| 215. | Sharp-shinned Hawk <i>Accipiter striatus</i> | 0.9288% |
| 216. | Cooper's Hawk <i>Accipiter cooperii</i> | 3.05% |
| 217. | Bald Eagle <i>Haliaeetus leucocephalus</i> | 12.81% |
| 218. | Red-shouldered Hawk <i>Buteo lineatus</i> | 25.33% |
| 219. | Broad-winged Hawk <i>Buteo platypterus</i> | 0.3852% |
| 220. | Short-tailed Hawk <i>Buteo brachyurus</i> | 0.9431% |
| 221. | Swainson's Hawk <i>Buteo swainsoni</i> | 0.0348% |
| 222. | Red-tailed Hawk <i>Buteo jamaicensis</i> | 3.33% |

| | | |
|------|--|----------------|
| 223. | Rough-legged Hawk <i>Buteo lagopus</i> | 0.0035% |
| 224. | Ferruginous Hawk <i>Buteo regalis</i> | 0.0000% |
| 225. | Barn Owl <i>Tyto alba</i> | 0.2233% |
| 226. | Eastern Screech-Owl <i>Megascops asio</i> | 0.9949% |
| 227. | Great Horned Owl <i>Bubo virginianus</i> | 2.02% |
| 228. | Burrowing Owl <i>Athene cunicularia</i> | 0.6832% |
| 229. | Barred Owl <i>Strix varia</i> | 4.66% |
| 230. | Long-eared Owl <i>Asio otus</i> | 0.0003% |
| 231. | Short-eared Owl <i>Asio flammeus</i> | 0.0185% |
| 232. | Northern Saw-whet Owl <i>Aegolius acadicus</i> | 0.0000% |
| 233. | Belted Kingfisher <i>Megaceryle alcyon</i> | 13.49% |
| 234. | Yellow-bellied Sapsucker <i>Sphyrapicus varius</i> | 3.85% |
| 235. | Red-headed Woodpecker <i>Melanerpes erythrocephalus</i> | 1.66% |
| 236. | Red-bellied Woodpecker <i>Melanerpes carolinus</i> | 42.08% |
| 237. | Downy Woodpecker <i>Dryobates pubescens</i> | 18.55% |
| 238. | Red-cockaded Woodpecker <i>Dryobates borealis</i> | 0.5947% |
| 239. | Hairy Woodpecker <i>Dryobates villosus</i> | 0.2811% |
| 240. | Ivory-billed Woodpecker <i>Campephilus principalis</i> | 0.0010% |
| 241. | Pileated Woodpecker <i>Dryocopus pileatus</i> | 15.86% |
| 242. | Northern Flicker <i>Colaptes auratus</i> | 3.97% |
| 243. | Crested Caracara <i>Caracara plancus</i> | 1.20% |
| 244. | Yellow-headed Caracara <i>Daptrius chimachima</i> | 0.0109% |

| | | |
|------|--|----------------|
| 245. | Eurasian Kestrel <i>Falco tinnunculus</i> | 0.0067% |
| 246. | American Kestrel <i>Falco sparverius</i> | 9.73% |
| 247. | Merlin <i>Falco columbarius</i> | 0.9055% |
| 248. | Peregrine Falcon <i>Falco peregrinus</i> | 0.7323% |
| 249. | Monk Parakeet <i>Myiopsitta monachus</i> | 1.62% |
| 250. | White-winged Parakeet <i>Brotogeris versicolurus</i> | 0.0524% |
| 251. | Carolina Parakeet <i>Conuropsis carolinensis</i> | 0.0000% |
| 252. | Nanday Parakeet <i>Aratinga nenday</i> | 2.87% |
| 253. | Mitred Parakeet <i>Psittacara mitratus</i> | 0.2328% |
| 254. | Eastern Wood-Pewee <i>Contopus virens</i> | 0.0029% |
| 255. | Cuban Pewee <i>Contopus caribaeus</i> | 0.0038% |
| 256. | Yellow-bellied Flycatcher <i>Empidonax flaviventris</i> | 0.0131% |
| 257. | Acadian Flycatcher <i>Empidonax virescens</i> | 0.0016% |
| 258. | Willow Flycatcher <i>Empidonax traillii</i> | 0.0019% |
| 259. | Least Flycatcher <i>Empidonax minimus</i> | 0.0699% |
| 260. | Hammond's Flycatcher <i>Empidonax hammondii</i> | 0.0038% |
| 261. | Western Flycatcher <i>Empidonax difficilis</i> | 0.0195% |
| 262. | Eastern Phoebe <i>Sayornis phoebe</i> | 10.86% |
| 263. | Say's Phoebe <i>Sayornis saya</i> | 0.0128% |
| 264. | Vermilion Flycatcher <i>Pyrocephalus rubinus</i> | 0.1198% |
| 265. | Ash-throated Flycatcher <i>Myiarchus cinerascens</i> | 0.0811% |
| 266. | Great Crested Flycatcher <i>Myiarchus crinitus</i> | 9.18% |

| | | |
|------|---|---------|
| 267. | Brown-crested Flycatcher <i>Myiarchus tyrannulus</i> | 0.0444% |
| 268. | La Sagra's Flycatcher <i>Myiarchus sagrae</i> | 0.0993% |
| 269. | Piratic Flycatcher <i>Legatus leucophaeus</i> | 0.0000% |
| 270. | Tropical Kingbird <i>Tyrannus melancholicus</i> | 0.1341% |
| 271. | Couch's Kingbird <i>Tyrannus couchii</i> | 0.0013% |
| 272. | Cassin's Kingbird <i>Tyrannus vociferans</i> | 0.0249% |
| 273. | Western Kingbird <i>Tyrannus verticalis</i> | 0.4155% |
| 274. | Eastern Kingbird <i>Tyrannus tyrannus</i> | 0.5308% |
| 275. | Gray Kingbird <i>Tyrannus dominicensis</i> | 0.1635% |
| 276. | Loggerhead Kingbird <i>Tyrannus caudifasciatus</i> | 0.0041% |
| 277. | Scissor-tailed Flycatcher <i>Tyrannus forficatus</i> | 0.3516% |
| 278. | Fork-tailed Flycatcher <i>Tyrannus savana</i> | 0.0019% |
| 279. | White-eyed Vireo <i>Vireo griseus</i> | 14.95% |
| 280. | Thick-billed Vireo <i>Vireo crassirostris</i> | 0.0652% |
| 281. | Bell's Vireo <i>Vireo bellii</i> | 0.0338% |
| 282. | Yellow-throated Vireo <i>Vireo flavifrons</i> | 2.51% |
| 283. | Blue-headed Vireo <i>Vireo solitarius</i> | 5.40% |
| 284. | Philadelphia Vireo <i>Vireo philadelphicus</i> | 0.0038% |
| 285. | Warbling Vireo <i>Vireo gilvus</i> | 0.0233% |
| 286. | Red-eyed Vireo <i>Vireo olivaceus</i> | 1.93% |
| 287. | Black-whiskered Vireo <i>Vireo altiloquus</i> | 0.0418% |
| 288. | Loggerhead Shrike <i>Lanius ludovicianus</i> | 7.24% |
| 289. | Blue Jay <i>Cyanocitta cristata</i> | 35.31% |

| | | |
|------|--|---------|
| 290. | Florida Scrub-Jay <i>Aphelocoma coerulescens</i> | 1.25% |
| 291. | American Crow <i>Corvus brachyrhynchos</i> | 18.40% |
| 292. | Fish Crow <i>Corvus ossifragus</i> | 36.42% |
| 293. | Carolina Chickadee <i>Poecile carolinensis</i> | 11.15% |
| 294. | Tufted Titmouse <i>Baeolophus bicolor</i> | 21.47% |
| 295. | Horned Lark <i>Eremophila alpestris</i> | 0.0083% |
| 296. | Bank Swallow <i>Riparia riparia</i> | 0.0185% |
| 297. | Tree Swallow <i>Tachycineta bicolor</i> | 10.89% |
| 298. | Bahama Swallow <i>Tachycineta cyaneoviridis</i> | 0.0010% |
| 299. | Purple Martin <i>Progne subis</i> | 6.66% |
| 300. | Northern Rough-winged Swallow <i>Stelgidopteryx serripennis</i> | 2.10% |
| 301. | Barn Swallow <i>Hirundo rustica</i> | 3.49% |
| 302. | Cliff Swallow <i>Petrochelidon pyrrhonota</i> | 0.0869% |
| 303. | Cave Swallow <i>Petrochelidon fulva</i> | 0.1543% |
| 304. | Red-whiskered Bulbul <i>Pycnonotus jocosus</i> | 0.1089% |
| 305. | Ruby-crowned Kinglet <i>Corthylio calendula</i> | 8.29% |
| 306. | Golden-crowned Kinglet <i>Regulus satrapa</i> | 0.0853% |
| 307. | White-breasted Nuthatch <i>Sitta carolinensis</i> | 0.3194% |
| 308. | Brown-headed Nuthatch <i>Sitta pusilla</i> | 2.07% |
| 309. | Red-breasted Nuthatch <i>Sitta canadensis</i> | 0.0632% |
| 310. | Brown Creeper <i>Certhia americana</i> | 0.0163% |
| 311. | Blue-gray Gnatcatcher <i>Polioptila caerulea</i> | 19.63% |

| | | |
|------|--|----------------|
| 312. | House Wren <i>Troglodytes aedon</i> | 6.94% |
| 313. | Winter Wren <i>Troglodytes hiemalis</i> | 0.0278% |
| 314. | Sedge Wren <i>Cistothorus stellaris</i> | 0.7934% |
| 315. | Marsh Wren <i>Cistothorus palustris</i> | 1.87% |
| 316. | Carolina Wren <i>Thryothorus ludovicianus</i> | 26.93% |
| 317. | Bewick's Wren <i>Thryomanes bewickii</i> | 0.0006% |
| 318. | European Starling <i>Sturnus vulgaris</i> | 12.55% |
| 319. | Common Myna <i>Acridotheres tristis</i> | 0.2782% |
| 320. | Gray Catbird <i>Dumetella carolinensis</i> | 27.43% |
| 321. | Pearly-eyed Thrasher <i>Margarops fuscatus</i> | 0.0006% |
| 322. | Brown Thrasher <i>Toxostoma rufum</i> | 8.46% |
| 323. | Sage Thrasher <i>Oreoscoptes montanus</i> | 0.0003% |
| 324. | Bahama Mockingbird <i>Mimus gundlachii</i> | 0.0297% |
| 325. | Northern Mockingbird <i>Mimus polyglottos</i> | 40.00% |
| 326. | Eastern Bluebird <i>Sialia sialis</i> | 10.11% |
| 327. | Mountain Bluebird <i>Sialia currucoides</i> | 0.0045% |
| 328. | Townsend's Solitaire <i>Myadestes townsendi</i> | 0.0006% |
| 329. | Veery <i>Catharus fuscescens</i> | 0.0013% |
| 330. | Gray-cheeked Thrush <i>Catharus minimus</i> | 0.0013% |
| 331. | Swainson's Thrush <i>Catharus ustulatus</i> | 0.0188% |
| 332. | Hermit Thrush <i>Catharus guttatus</i> | 2.30% |
| 333. | Wood Thrush <i>Hylocichla mustelina</i> | 0.0936% |
| 334. | American Robin <i>Turdus migratorius</i> | 4.72% |

| | | |
|------|---|---------|
| 335. | Red-legged Thrush <i>Turdus plumbeus</i> | 0.0080% |
| 336. | Cedar Waxwing <i>Bombycilla cedrorum</i> | 7.50% |
| 337. | Tricolored Munia <i>Lonchura malacca</i> | 0.0355% |
| 338. | House Sparrow <i>Passer domesticus</i> | 4.89% |
| 339. | White Wagtail <i>Motacilla alba</i> | 0.0013% |
| 340. | American Pipit <i>Anthus rubescens</i> | 0.1348% |
| 341. | Sprague's Pipit <i>Anthus spragueii</i> | 0.0019% |
| 342. | Evening Grosbeak <i>Coccothraustes vespertinus</i> | 0.0003% |
| 343. | House Finch <i>Haemorhous mexicanus</i> | 5.59% |
| 344. | Purple Finch <i>Haemorhous purpureus</i> | 0.0747% |
| 345. | Pine Siskin <i>Spinus pinus</i> | 0.3130% |
| 346. | American Goldfinch <i>Spinus tristis</i> | 6.87% |
| 347. | Lapland Longspur <i>Calcarius lapponicus</i> | 0.0013% |
| 348. | Snow Bunting <i>Plectrophenax nivalis</i> | 0.0013% |
| 349. | Little Bunting <i>Emberiza pusilla</i> | 0.0182% |
| 350. | Bachman's Sparrow <i>Peucaea aestivalis</i> | 1.17% |
| 351. | Grasshopper Sparrow <i>Ammodramus savannarum</i> | 0.4842% |
| 352. | Chipping Sparrow <i>Spizella passerina</i> | 6.20% |
| 353. | Clay-colored Sparrow <i>Spizella pallida</i> | 0.1000% |
| 354. | Field Sparrow <i>Spizella pusilla</i> | 0.2654% |
| 355. | Lark Sparrow <i>Chondestes grammacus</i> | 0.0610% |
| 356. | Lark Bunting <i>Calamospiza melanocorys</i> | 0.0010% |
| 357. | Fox Sparrow <i>Passerella iliaca</i> | 0.0038% |

| | | |
|------|---|---------|
| 358. | Dark-eyed Junco <i>Junco hyemalis</i> | 0.0131% |
| 359. | White-crowned Sparrow <i>Zonotrichia leucophrys</i> | 0.3513% |
| 360. | Harris's Sparrow <i>Zonotrichia querula</i> | 0.0144% |
| 361. | White-throated Sparrow <i>Zonotrichia albicollis</i> | 0.6254% |
| 362. | Vesper Sparrow <i>Pooecetes gramineus</i> | 0.3258% |
| 363. | LeConte's Sparrow <i>Ammospiza leconteii</i> | 0.0230% |
| 364. | Seaside Sparrow <i>Ammospiza maritima</i> | 0.1626% |
| 365. | Nelson's Sparrow <i>Ammospiza nelsoni</i> | 0.0578% |
| 366. | Saltmarsh Sparrow <i>Ammospiza caudacuta</i> | 0.0444% |
| 367. | Savannah Sparrow <i>Passerculus sandwichensis</i> | 4.81% |
| 368. | Henslow's Sparrow <i>Centronyx henslowii</i> | 0.0434% |
| 369. | Song Sparrow <i>Melospiza melodia</i> | 1.12% |
| 370. | Lincoln's Sparrow <i>Melospiza lincolnii</i> | 0.0722% |
| 371. | Swamp Sparrow <i>Melospiza georgiana</i> | 3.57% |
| 372. | Green-tailed Towhee <i>Pipilo chlorurus</i> | 0.0163% |
| 373. | Spotted Towhee <i>Pipilo maculatus</i> | 0.0109% |
| 374. | Eastern Towhee <i>Pipilo erythrophthalmus</i> | 7.11% |
| 375. | Western Spindalis <i>Spindalis zena</i> | 0.0760% |
| 376. | Yellow-breasted Chat <i>Icteria virens</i> | 0.0821% |
| 377. | Yellow-headed Blackbird <i>Xanthocephalus xanthocephalus</i> | 0.1223% |
| 378. | Bobolink <i>Dolichonyx oryzivorus</i> | 0.0051% |
| 379. | Western Meadowlark <i>Sturnella neglecta</i> | 0.0029% |

| | | |
|------|--|----------------|
| 380. | Eastern Meadowlark <i>Sturnella magna</i> | 4.61% |
| 381. | Orchard Oriole <i>Icterus spurius</i> | 0.1246% |
| 382. | Bullock's Oriole <i>Icterus bullockii</i> | 0.0198% |
| 383. | Spot-breasted Oriole <i>Icterus pectoralis</i> | 0.2718% |
| 384. | Baltimore Oriole <i>Icterus galbula</i> | 1.08% |
| 385. | Scott's Oriole <i>Icterus parisorum</i> | 0.0067% |
| 386. | Red-winged Blackbird <i>Agelaius phoeniceus</i> | 26.49% |
| 387. | Shiny Cowbird <i>Molothrus bonariensis</i> | 0.0882% |
| 388. | Bronzed Cowbird <i>Molothrus aeneus</i> | 0.3242% |
| 389. | Brown-headed Cowbird <i>Molothrus ater</i> | 6.13% |
| 390. | Rusty Blackbird <i>Euphagus carolinus</i> | 0.0460% |
| 391. | Brewer's Blackbird <i>Euphagus cyanocephalus</i> | 0.0176% |
| 392. | Common Grackle <i>Quiscalus quiscula</i> | 24.86% |
| 393. | Boat-tailed Grackle <i>Quiscalus major</i> | 29.84% |
| 394. | Great-tailed Grackle <i>Quiscalus mexicanus</i> | 0.0300% |
| 395. | Ovenbird <i>Seiurus aurocapilla</i> | 1.84% |
| 396. | Worm-eating Warbler <i>Helmitheros vermivorum</i> | 0.3293% |
| 397. | Louisiana Waterthrush <i>Parkesia motacilla</i> | 0.7512% |
| 398. | Northern Waterthrush <i>Parkesia noveboracensis</i> | 0.9872% |
| 399. | Bachman's Warbler <i>Vermivora bachmanii</i> | 0.0000% |
| 400. | Golden-winged Warbler <i>Vermivora chrysoptera</i> | 0.0019% |
| 401. | Blue-winged Warbler <i>Vermivora cyanoptera</i> | 0.0377% |
| 402. | Black-and-white Warbler <i>Mniotilta varia</i> | 9.06% |

| | | |
|------|--|---------|
| 403. | Prothonotary Warbler <i>Protonotaria citrea</i> | 0.5666% |
| 404. | Swainson's Warbler <i>Limnothlypis swainsonii</i> | 0.1217% |
| 405. | Tennessee Warbler <i>Leiothlypis peregrina</i> | 0.0169% |
| 406. | Orange-crowned Warbler <i>Leiothlypis celata</i> | 1.72% |
| 407. | Nashville Warbler <i>Leiothlypis ruficapilla</i> | 0.0278% |
| 408. | Connecticut Warbler <i>Oporornis agilis</i> | 0.0003% |
| 409. | MacGillivray's Warbler <i>Geothlypis tolmiei</i> | 0.0125% |
| 410. | Kentucky Warbler <i>Geothlypis formosa</i> | 0.0319% |
| 411. | Common Yellowthroat <i>Geothlypis trichas</i> | 11.20% |
| 412. | Hooded Warbler <i>Setophaga citrina</i> | 0.8320% |
| 413. | American Redstart <i>Setophaga ruticilla</i> | 1.20% |
| 414. | Cape May Warbler <i>Setophaga tigrina</i> | 0.2692% |
| 415. | Cerulean Warbler <i>Setophaga cerulea</i> | 0.0013% |
| 416. | Northern Parula <i>Setophaga americana</i> | 24.98% |
| 417. | Magnolia Warbler <i>Setophaga magnolia</i> | 0.3092% |
| 418. | Bay-breasted Warbler <i>Setophaga castanea</i> | 0.0006% |
| 419. | Blackburnian Warbler <i>Setophaga fusca</i> | 0.0038% |
| 420. | Yellow Warbler <i>Setophaga petechia</i> | 0.1530% |
| 421. | Chestnut-sided Warbler <i>Setophaga pensylvanica</i> | 0.0077% |
| 422. | Blackpoll Warbler <i>Setophaga striata</i> | 0.0038% |
| 423. | Black-throated Blue Warbler <i>Setophaga caerulescens</i> | 0.3389% |
| 424. | Palm Warbler <i>Setophaga palmarum</i> | 33.51% |

| | |
|---|----------------|
| 425. Pine Warbler <i>Setophaga pinus</i> | 11.77% |
| 426. Yellow-rumped Warbler <i>Setophaga coronata</i> | 32.13% |
| 427. Yellow-throated Warbler <i>Setophaga dominica</i> | 5.42% |
| 428. Prairie Warbler <i>Setophaga discolor</i> | 6.70% |
| 429. Black-throated Gray Warbler <i>Setophaga nigrescens</i> | 0.0067% |
| 430. Townsend's Warbler <i>Setophaga townsendi</i> | 0.0141% |
| 431. Black-throated Green Warbler <i>Setophaga virens</i> | 0.6822% |
| 432. Wilson's Warbler <i>Cardellina pusilla</i> | 0.0843% |
| 433. Summer Tanager <i>Piranga rubra</i> | 0.8368% |
| 434. Scarlet Tanager <i>Piranga olivacea</i> | 0.0086% |
| 435. Western Tanager <i>Piranga ludoviciana</i> | 0.1999% |
| 436. Northern Cardinal <i>Cardinalis cardinalis</i> | 52.66% |
| 437. Rose-breasted Grosbeak <i>Pheucticus ludovicianus</i> | 0.0731% |
| 438. Black-headed Grosbeak <i>Pheucticus melanocephalus</i> | 0.0041% |
| 439. Blue Grosbeak <i>Passerina caerulea</i> | 0.1201% |
| 440. Lazuli Bunting <i>Passerina amoena</i> | 0.0310% |
| 441. Indigo Bunting <i>Passerina cyanea</i> | 1.80% |
| 442. Painted Bunting <i>Passerina ciris</i> | 5.58% |
| 443. Dickcissel <i>Spiza americana</i> | 0.0530% |
| 444. Red-legged Honeycreeper <i>Cyanerpes cyaneus</i> | 0.0089% |
| 445. Bananaquit <i>Coereba flaveola</i> | 0.0074% |
| 446. Yellow-faced Grassquit <i>Tiaris olivaceus</i> | 0.0016% |

447. **Black-faced Grassquit** *Melanospiza bicolor* **0.0460%**

EXOTIC: PROVISIONAL

448. **Indian Peafowl** *Pavo cristatus* **0.4762%** frequency

449. **Red Junglefowl** *Gallus gallus* **0.1380%**

450. **Red-footed Booby** *Sula sula* **0.0022%**

451. **Great White Pelican** *Pelecanus onocrotalus* **0.0032%**

452. **Scarlet Ibis** *Eudocimus ruber* **0.0000%**

453. **Rose-ringed Parakeet** *Psittacula krameri* **0.0355%**

454. **Budgerigar** *Melopsittacus undulatus* **0.0077%**

455. **Yellow-chevroned Parakeet** *Brotogeris chiriri* **0.2044%**

456. **Red-crowned Parrot** *Amazona viridigenalis* **0.0600%**

457. **Orange-winged Parrot** *Amazona amazonica* **0.1236%**

458. **Blue-and-yellow Macaw** *Ara ararauna* **0.0757%**

459. **Chestnut-fronted Macaw** *Ara severus* **0.0814%**

460. **Blue-crowned Parakeet** *Thectocercus acuticaudatus* **0.3593%**

461. **Red-masked Parakeet** *Psittacara erythrogenys* **0.1750%**

462. **White-eyed Parakeet** *Psittacara leucophthalmus* **0.0479%**

463. **House Crow** *Corvus splendens* **0.0188%**

464. **Common Hill Myna** *Gracula religiosa* **0.0661%**

465. **Scaly-breasted Munia** *Lonchura punctulata* **0.2287%**

EXOTIC: ESCAPEE

Plumed Whistling-Duck *Dendrocygna eytoni* **0.0003%** frequency

| | |
|--|----------------|
| White-faced Whistling-Duck <i>Dendrocygna viduata</i> | 0.0032% |
| Bar-headed Goose <i>Anser indicus</i> | 0.0003% |
| Graylag Goose <i>Anser anser</i> | 0.1172% |
| Swan Goose <i>Anser cygnoides</i> | 0.1000% |
| Barnacle Goose <i>Branta leucopsis</i> | 0.0000% |
| Blue-winged Goose <i>Cyanochen cyanoptera</i> | 0.0003% |
| Mute Swan <i>Cygnus olor</i> | 0.2421% |
| Black Swan <i>Cygnus atratus</i> | 0.0910% |
| Black-necked Swan <i>Cygnus melancoryphus</i> | 0.0262% |
| Trumpeter Swan <i>Cygnus buccinator</i> | 0.0294% |
| Whooper Swan <i>Cygnus cygnus</i> | 0.0153% |
| Coscoroba Swan <i>Coscoroba coscoroba</i> | 0.0048% |
| Radjah Shelduck <i>Radjah radjah</i> | 0.0000% |
| Ruddy Shelduck <i>Tadorna ferruginea</i> | 0.0086% |
| Common Shelduck <i>Tadorna tadorna</i> | 0.0099% |
| Ringed Teal <i>Callonetta leucophrys</i> | 0.0131% |
| Mandarin Duck <i>Aix galericulata</i> | 0.0112% |
| Chiloe Wigeon <i>Mareca sibilatrix</i> | 0.0013% |
| Red-crested Pochard <i>Netta rufina</i> | 0.0019% |
| Rosy-billed Pochard <i>Netta peposaca</i> | 0.0010% |
| Helmeted Guineafowl <i>Numida meleagris</i> | 0.0252% |
| Ring-necked Pheasant <i>Phasianus colchicus</i> | 0.0083% |

| | |
|--|---------|
| Green Pheasant <i>Phasianus versicolor</i> | 0.0000% |
| Green Peafowl <i>Pavo muticus</i> | 0.0019% |
| Red-legged Partridge <i>Alectoris rufa</i> | 0.0029% |
| African Collared-Dove <i>Streptopelia roseogrisea</i> | 0.0016% |
| Diamond Dove <i>Geopelia cuneata</i> | 0.0003% |
| White-cheeked Turaco <i>Menelikornis leucotis</i> | 0.0000% |
| Black-necked Stork <i>Ephippiorhynchus asiaticus</i> | 0.0003% |
| African Sacred Ibis <i>Threskiornis aethiopicus</i> | 0.0000% |
| Great Black Hawk <i>Buteogallus urubitinga</i> | 0.0029% |
| Black-casqued Hornbill <i>Ceratogymna atrata</i> | 0.0003% |
| Cockatiel <i>Nymphicus hollandicus</i> | 0.0051% |
| Red-breasted Parakeet <i>Psittacula alexandri</i> | 0.0003% |
| Rosy-faced Lovebird <i>Agapornis roseicollis</i> | 0.0010% |
| Yellow-collared Lovebird <i>Agapornis personatus</i> | 0.0003% |
| Scaly-headed Parrot <i>Pionus maximiliani</i> | 0.0038% |
| Lilac-crowned Parrot <i>Amazona finschi</i> | 0.0096% |
| Red-lored Parrot <i>Amazona autumnalis</i> | 0.0003% |
| Yellow-naped Parrot <i>Amazona auropalliata</i> | 0.0006% |
| Yellow-headed Parrot <i>Amazona oratrix</i> | 0.0019% |
| Yellow-crowned Parrot <i>Amazona ochrocephala</i> | 0.0112% |
| Turquoise-fronted Parrot <i>Amazona aestiva</i> | 0.0086% |
| White-fronted Parrot <i>Amazona albifrons</i> | 0.0019% |

| | |
|---|---------|
| Mealy Parrot <i>Amazona farinosa</i> | 0.0006% |
| Green-cheeked Parakeet <i>Pyrrhura molinae</i> | 0.0000% |
| Peach-fronted Parakeet <i>Eupsittula aurea</i> | 0.0000% |
| Dusky-headed Parakeet <i>Aratinga weddellii</i> | 0.0000% |
| Sun Parakeet <i>Aratinga solstitialis</i> | 0.0000% |
| Military Macaw <i>Ara militaris</i> | 0.0000% |
| Scarlet Macaw <i>Ara macao</i> | 0.0000% |
| Red-shouldered Macaw <i>Diopsittaca nobilis</i> | 0.0000% |
| Green Parakeet <i>Psittacara holochlorus</i> | 0.0105% |
| Crimson-fronted Parakeet <i>Psittacara finschi</i> | 0.0118% |
| Crested Myna <i>Acridotheres cristatellus</i> | 0.0000% |
| Tropical Mockingbird <i>Mimus gilvus</i> | 0.0041% |
| Northern Red Bishop <i>Euplectes franciscanus</i> | 0.0003% |
| Zebra Finch <i>Taeniopygia guttata</i> | 0.0070% |
| Java Sparrow <i>Padda oryzivora</i> | 0.0000% |
| Yellow-fronted Canary <i>Crithagra mozambica</i> | 0.0003% |
| Venezuelan Troupial <i>Icterus icterus</i> | 0.0000% |
| Red-crested Cardinal <i>Paroaria coronata</i> | 0.0000% |
| Paradise Tanager <i>Tangara chilensis</i> | 0.0003% |
| Saffron Finch <i>Sicalis flaveola</i> | 0.0003% |
| Cuban Grassquit <i>Phonipara canora</i> | 0.0003% |

HYBRIDS

| | |
|---|-------------------|
| Snow x Ross's Goose (hybrid) <i>Anser caerulescens x rossii</i> | 0.0003% frequency |
| Graylag x Swan Goose (hybrid) <i>Anser anser x cygnoides</i> | 0.0093% |
| Swan Goose x Canada Goose (hybrid) <i>Anser cygnoides x Branta canadensis</i> | 0.0013% |
| Domestic goose sp. x Canada Goose (hybrid) <i>Anser sp. (Domestic type) x Branta canadensis</i> | 0.0010% |
| Blue-winged x Cinnamon Teal (hybrid) <i>Spatula discors x cyanoptera</i> | 0.0172% |
| Blue-winged Teal x Northern Shoveler (hybrid) <i>Spatula discors x clypeata</i> | 0.0003% |
| Muscovy Duck x Mallard (hybrid) <i>Cairina moschata x Anas platyrhynchos</i> | 0.0265% |
| Mallard x Mottled Duck (hybrid) <i>Anas platyrhynchos x fulvigula</i> | 2.08% |
| American Wigeon x Northern Pintail (hybrid) <i>Mareca americana x Anas acuta</i> | 0.0003% |
| Redhead x Ring-necked Duck (hybrid) <i>Aythya americana x collaris</i> | 0.0067% |
| Tufted Duck x Lesser Scaup (hybrid) <i>Aythya fuligula x affinis</i> | 0.0006% |
| Redhead x scaup sp. (hybrid) <i>Aythya americana x marila/affinis</i> | 0.0010% |
| Eurasian Collared-Dove x Mourning Dove (hybrid) <i>Streptopelia decaocto x Zenaida macroura</i> | 0.0003% |
| Rufous x Allen's Hummingbird (hybrid) <i>Selasphorus rufus x sasin</i> | 0.0006% |
| Laughing x Ring-billed Gull (hybrid) <i>Leucophaeus atricilla x Larus delawarensis</i> | 0.0003% |
| Herring x Glaucous Gull (hybrid) <i>Larus argentatus x hyperboreus</i> | 0.0038% |

| | |
|--|----------------|
| Herring x Lesser Black-backed Gull (hybrid) <i>Larus argentatus x fuscus</i> | 0.0003% |
| Double-crested x Neotropic Cormorant (hybrid) <i>Nannopterum auritum x brasilianum</i> | 0.0032% |
| Little Blue Heron x Snowy Egret (hybrid) <i>Egretta caerulea x thula</i> | 0.0013% |
| Tricolored Heron x Snowy Egret (hybrid) <i>Egretta tricolor x thula</i> | 0.0080% |
| Great Egret x Great Blue Heron (hybrid) <i>Ardea alba x herodias</i> | 0.0064% |
| White x Scarlet Ibis (hybrid) <i>Eudocimus albus x ruber</i> | 0.0003% |
| Glossy x White-faced Ibis (hybrid) <i>Plegadis falcinellus x chihi</i> | 0.0006% |
| White-winged x Yellow-chevroned Parakeet (hybrid) <i>Brotogeris versicolurus x chiriri</i> | 0.0010% |
| House x Fish Crow (hybrid) <i>Corvus splendens x ossifragus</i> | 0.0003% |